

Bioanalytical characterisation of multiple endocrine- and dioxin-like activities in sediments from reference and impacted small rivers.

Said Kinani, Stéphane Bouchonnet, Nicolas Creusot, Sophie Bourcier, Patrick Balaguer, Jean-Marc Porcher, Sélim Aït-Aïssa

► **To cite this version:**

Said Kinani, Stéphane Bouchonnet, Nicolas Creusot, Sophie Bourcier, Patrick Balaguer, et al.. Bio-analytical characterisation of multiple endocrine- and dioxin-like activities in sediments from reference and impacted small rivers.. Environmental Pollution, Elsevier, 2010, 158 (1), pp.74-83. 10.1016/j.envpol.2009.07.041 . ineris-00453475

HAL Id: ineris-00453475

<https://hal-ineris.archives-ouvertes.fr/ineris-00453475>

Submitted on 4 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Bioanalytical characterisation of multiple endocrine- and dioxin-like activities in**
2 **sediments from reference and impacted small rivers**

3

4 Said Kinani^{a,b,*}, Stéphane Bouchonnet^{b,*}, Nicolas Creusot^a, Sophie Bourcier^b, Patrick
5 Balaguer^c, Jean-Marc Porcher^a and Sélim Aït-Aïssa^{a,1,*}

6

7 ^a Unité d'Ecotoxicologie, Institut National de l'Environnement Industriel et des Risques
8 (INERIS), BP2, F-60550 Verneuil en Halatte, France.

9 ^b Département de Chimie des Mécanismes Réactionnels, Ecole Polytechnique, 91128
10 Palaiseau Cedex, France.

11 ^c Institut National de la Santé et de la Recherche Médicale (INSERM), U896, Montpellier, F-
12 34298, France.

13

14 1 : Corresponding author: Tel +33 344 556 511, Fax +33 344 556 767

15 * E-mail addresses : selim.ait-aïssa@ineris.fr (Sélim Aït-Aïssa), said@dcmr.polytechnique.fr
16 (Said Kinani), stephane.bouchonnet@dcmr.polytechnique.fr (Stéphane Bouchonnet).

17

18 **Abstract**

19 A comprehensive evaluation of organic contamination was performed in sediments sampled
20 in two reference and three impacted small streams where endocrine disruptive (ED) effects in
21 fish have been evidenced. The approach combined quantitative chemical analyses of more
22 than 50 ED chemicals and *in vitro* bioassays allowing the quantification of receptor-mediated
23 activities, namely estrogen (ER), androgen (AR), dioxin (AhR) and pregnane X (PXR)
24 receptors. At the most impacted sites, chemical analyses showed the presence of natural
25 estrogens, organochlorine pesticides, parabens, polycyclic aromatic hydrocarbons (16 PAHs),
26 bisphenol A and alkylphenols, while synthetic steroids, myco-estrogens and phyto-estrogens
27 were not detected. Determination of toxic-equivalent amounts showed that 28 to 96% of
28 estrogenic activities in bioassays (0.2-6.3 ng/g 17 β -estradiol equivalents) were explained by
29 17 β -estradiol and estrone. PAHs were major contributors (20-60%) to the total dioxin-like
30 activities. Interestingly, high PXR and (anti)AR activities were detected; however, analysed
31 compounds could not explain the measured biological activities.

32 *Keywords:* river sediment; endocrine disrupting chemicals; *in vitro* bioassays; GC-MS & LC-
33 MS analysis; mass balance analysis

34 *Capsule :* Multiple endocrine disrupting chemicals (ER, AR, AhR and PXR ligands) are
35 detected in French river sediments using a panel of *in vitro* bioassays and analytical methods.

36

37 **1. Introduction**

38 The rapid industrial and urban development which occurred in the second half of the 20th
39 century allowed the emergence of millions of persistent anthropogenic chemicals in our
40 environment without prior study of their toxicity. In recent years, a newly defined category of
41 these chemicals, with the potential to interact with the endocrine system have emerged and
42 aroused considerable interest of scientific communities. Named endocrine disrupting-
43 chemicals (EDCs), these substances are defined as "exogenous agents that interfere with the
44 synthesis, secretion, transport, binding, action, or elimination of natural hormones which are
45 responsible for maintenance of homeostasis, reproduction, development and/or behavior"
46 (Kavlock et al., 1996). In the aquatic environment, occurring EDCs are very diverse in terms
47 of chemical nature and origins (Vos et al., 2000).

48 Up to date, most attention has been directed to identifying estrogenic chemicals, i.e. those
49 capable of binding to and activating the estrogen receptor (ER), because many of the effects
50 reported in wildlife appear to be a consequence of feminization of males (Sumpter et al.,
51 2005). However, many environmental contaminants can interfere with other molecular targets
52 involved in the regulation of the endocrine system. These include other nuclear receptors like
53 androgen receptor (AR), pregnane X receptor (PXR, e.g. Mnif et al., 2007) as well as
54 steroidogenesis enzymes like aromatase (e.g. Laville et al., 2006). EDCs can also act via
55 indirect mechanisms such as (anti)estrogenic effect mediated by aryl hydrocarbon receptor
56 (AhR) ligands through an ER/AhR cross-talk dependent mechanism (Othake et al., 2003,
57 2007). Therefore, a comprehensive assessment of EDC hazard should take into account the
58 ability of chemicals to interfere with different targets of the endocrine system (Houtman et al.,
59 2004).

60 Regarding the large diversity of EDCs and their effects, bioanalytical approaches using
61 mechanism-based biological screening tools have emerged to monitor such substances in

62 complex environmental mixtures (Eggen and Segner, 2003). In particular, *in vitro* assays
63 using reporter gene activation in stably transfected cell lines provide robust, sensitive and
64 specific bioassays to screen and quantify endocrine activities in environmental samples.
65 Combining such tools with powerful chemical analyses within integrated approaches permits
66 to identify key toxicants to be monitored in the environment (Brack, 2003). By using such
67 bioanalytical approaches, the occurrence of EDCs and their effects to aquatic organisms has
68 been several times reported in different industrialised countries like the United Kingdom (e.g.
69 Jobling et al., 2006) or the Netherlands (e.g. Vethaak et al., 2005). However in France, much
70 fewer data have been reported. Recent studies showed occurrence of steroid estrogens and
71 alkylphenols in effluents from sewage treatment plants (Labadie and Budzinski, 2005, Muller
72 et al., 2008) or in water and sediment from the Seine River (Fenet et al., 2003; Cargouet et al.,
73 2004). More recently, abnormal elevated levels of vitellogenin and spigging, used as
74 biomarkers of estrogen and androgen exposure respectively, have been reported in three-spine
75 stickleback (*Gasterosteus aculeatus*) sampled in different small streams subjected to diffuse
76 pollution like agricultural run-off or domestic effluents (Sanchez et al., 2008). Thus, more
77 investigation is needed in order to characterise and identify key EDCs in French aquatic
78 ecosystems.

79 This study aimed at performing a comprehensive evaluation of the chemical contamination by
80 EDCs in sediments sampled in small streams at sites where endocrine disruptive effects in
81 fish have been previously evidenced (Sanchez et al., 2007, Sanchez et al., 2008). The five
82 selected sites, located in the North of France, were representative of different levels of water
83 and ecological quality (Table 1). For this purpose, we used a combined approach involving i)
84 targeted chemical analyses of more than 50 chemicals (Table 2) selected on the basis of both
85 their environmental occurrence and their known EDC potency, and ii) a panel of *in vitro*
86 bioassays that allowed the detection of different receptor-mediated activities, namely ER, AR,

87 AhR and PXR. Since all activities were detected in at least two of the five studied sites, the
88 contribution of analyzed EDCs in the biological activities detected by the bioassays was
89 estimated by comparing toxic-equivalent quantities from both approaches.

90

91 **2. Materials and methods**

92

93 **2.1. Chemicals and reagents**

94 List, abbreviation and source of analytical standards are given in Table 2. Flutamide (Flut), 3-
95 (4,5-dimethylthiazol-2-ol)-2,5-diphenyltetrasodium bromide tetrazolium (MTT), 5 α -
96 dihydrotestosterone (DHT), as well as the analytical internal standards including [¹³C₆]4-n-
97 nonylphenol, [²H₂]17 β -estradiol, [²H₁₀]phenanthrene, [²H₁₀]acenaphthene, [²H₁₂]perylene and
98 [²H₁₂]chrysene were purchased from Sigma-Aldrich (St Quentin Fallavier, France). ICI
99 182,780 (ICI) and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) were obtained from Tocris
100 Bioscience (Ellisville, USA) and Promochem (Molsheim, France), respectively. All standards
101 were of 98.1–99.9% purity. Chromatographic-grade solvents: acetone, methanol, acetonitrile,
102 dimethylsulfoxide (DMSO) and n-hexane were also purchased from Sigma-Aldrich. Stock
103 solutions of β -E2 2.72 g/L, DHT 2.90 g/L and B[a]P 2.52 g/L were prepared by dissolving
104 crystalline compound in DMSO. A solution of TCDD was prepared by drying an aliquot of
105 TCDD in methanol under a nitrogen stream and dissolving the residue in DMSO to reach a
106 concentration of 3.22 g/L.

107

108 **2.2. Site description and sampling**

109 Surface sediment samples were collected in five rivers located in the North of France during
110 July 2004. The selected stations presented different levels of impacts in terms of
111 anthropogenic pressures, chemical contamination and impacts on fish populations (Table 1):

112 the Aisne (Ais) and the Vallon du Vivier (VdV) rivers were considered as reference sites; the
113 Lézarde (Lez) as a moderately impacted site and the Réveillon (Rév) and Rhonelle (Rho) as
114 heavily impacted sites. At each site, at least five grab sediment samples were taken alongside
115 the river section of the study sites and were pooled to obtain an average sediment sample and
116 to minimize intra-site variability. All sediment samples were passed through a 2 mm sieve
117 pore, stored in aluminium boxes and immediately transferred to the laboratory where they
118 were stored at -20 °C in order to inhibit biological activity until extraction.

119

120 ***2.3. Extraction procedure***

121 The sediment samples were extracted as previously described (Kinani et al., 2008a). Briefly, 5
122 g of lyophilised and homogenised sediment were extracted three times with 10 mL of a
123 hexane/acetone (2:1, v/v) mixture. For each extraction step, the sample was sonicated 10 min
124 (ENMA Transsonic 460/H, Frequency 50-60 Hz, Germany) at room temperature. The extracts
125 were combined and the supernatant was transferred to a 50 mL vial containing 1.0 g of
126 anhydrous sodium sulphate. The organic extracts were then centrifuged for 10 min at 3000
127 rpm and the supernatant was collected, filtered onto a 0.45 µm Acrodisc CRPTFE Syringe
128 filter (GelmanSciences, USA), reduced to about 1 mL using rotary evaporation at 30 °C or
129 lower, evaporated to complete dryness under a gentle nitrogen stream and reconstituted into 1
130 mL of the extraction solvent. The final extract was then divided into two parts, the first one
131 for chemical analysis and the second one for bioassay testing. The extraction solvent was
132 subsequently replaced by DMSO for bioassay experiments.

133

134 ***2.4. Cell cultures and in vitro bioassays***

135 ***2.4.1. Estrogenic, PXR and (anti-)androgenic activities in reporter gene assays***

136 The estrogenic, PXR and (anti-)androgenic activities of the extracts were monitored by using
137 the MELN, HG₅LN-PXR and MDA-kb2 reporter cell lines, respectively. The MELN cell line
138 was obtained by stable transfection of MCF-7 human breast cancer cells by an ERE-βGlob-
139 Luc-SVNeo plasmid (Balaguer et al., 2001). The HG₅LN-PXR cell line was derived from
140 HeLa cells that were first stably transfected with GAL4RE₅-bGlob-Luc-SV-Neo (HG₅LN
141 cells) before being stably transfected with the pSG5-GAL4(DBD)-hPXR(LBD)-puro plasmid
142 (Lemaire et al., 2006). The MDA-kb2 cell line (ATCC, #CRL-2713) was derived from the
143 MDA-MB-453 human breast cancer cells. They were stably transfected by a MMTV
144 promoter-luciferase plasmid construct, which is under the control of endogenous AR and
145 glucocorticoid receptor (GR) (Wilson et al., 2002). All reporter cell lines were routinely
146 cultured in phenol red containing Dulbecco's Modified Eagle's Medium (DMEM),
147 supplemented with 5% foetal calf serum (FCS), 1% nonessential amino acids and
148 penicillin/streptomycin (50 U/mL each) in a 5% CO₂ humidified atmosphere at 37 °C. For
149 experiments, cells were left to incubate for two days in phenol red free DMEM supplemented
150 with 3% dextran charcoal coated-FCS (DCC medium) before seeded in white opaque 96-
151 wells culture plates at a density of 50 000 cells per well. Serial dilutions of reference
152 chemicals or organic extracts were added in triplicates 24 h later and then left to incubate for
153 16 h. In all assays, DMSO in the culture medium was always at 0.5% v/v, including in cell
154 controls. At this concentration, it did not affect cell viability or luciferase activity. At the end
155 of the incubation period, the medium was removed and replaced by 50 µL of DCC medium
156 containing 0.3 mM of D-luciferin and the luminescence signal was measured in living cells
157 for 2 seconds per well with a microtiter plate luminometer (µBeta, Wallac). Relative
158 luminescence units (RLU) were converted to relative response units expressed as percent of
159 maximal luciferase activity induced by the positive controls: E2 10 nM, SR12813 0.3 µM,
160 DHT 10 nM (for AR agonist activity) and DHT 0.3 nM (for AR antagonist activity) in

161 MELN, HG₅LN-PXR and MDA-kb2 cell lines, respectively. For ER and AR agonistic
162 responses, the specificity of luciferase induction by samples was checked by co-exposure
163 experiments with 0.1 μM ICI182,780 or 1 μM flutamide, as reference ER and AR antagonists,
164 respectively.

165 *2.4.2. Dioxin-like activity in PLHC-1 cell line*

166 The fish hepatic PLHC-1 cell line (ATCC, #CRL-2406) was described by Ryan and
167 Hightower (1994). The cells were routinely grown at 30 °C in E-MEM culture media
168 supplemented with 10 % FCS and 1 % antibiotics in a 5 % CO₂ humidified atmosphere. For
169 experiments, cells were seeded in 96-well plates at a density of approximately 50 000 cells
170 per well. After 24 hours of incubation, cells were exposed to test chemicals or sample extracts
171 for either a 4 h or 24 h incubation period, in order to differentiate between dioxin-like
172 compounds that are rapidly metabolized (e.g. PAHs) in the cells and dioxin-like chemicals
173 that are persistent in the cells (e.g. dioxins) (Louiz et al., 2008). Then, plates were processed
174 for 7-ethoxyresorufin-O-deethylase (EROD) activity in intact cells, following the protocol
175 previously described by Laville et al. (2004). Results were expressed as percent of EROD
176 activity induced by the positive control (TCDD 1 nM).

177 *2.4.3. Cell viability*

178 The effect of test compounds or samples on cellular viability in the different cell lines was
179 evaluated by using the methyl-thiazol-tetrazolium (MTT) assay (Mosmann, 1983), as
180 previously described (Laville et al., 2004).

181

182 *2.5. Chemical analyses*

183 Two mass spectrometers coupled with chromatography have been used for the quantitative
184 analysis: a “CP3800” gas chromatograph system equipped with a “CP8400” autosampler and
185 coupled to a “Saturn 2000” ion trap mass spectrometer (Varian, Les Ulis, France) for the GC-

186 MS and a “alliance 2690” liquid chromatography coupled to a Q-TOF Premier (Waters,
187 France) mass spectrometer for LC-MS/MS. Before performing the chemical analysis, a
188 portion of each extract was purified according to a method previously described by Hartmann
189 et al., (2007).

190 2.5.1. GC-MS and GC-MS/MS analyses

191 The chromatographic separation was performed with a 60 m “Factor four VF-10-MS” (10%
192 phenyl, 90 % methylpolysiloxane) capillary column (internal diameter: 0.25 mm, film
193 thickness: 0.25 μ m) from Varian. All experiments were performed by automatically injecting
194 2.0 μ L of sample in the splitless mode at a rate of 50 μ L/s. Helium (99.999% purity) was used
195 as the carrier gas at a constant flow of 1.0 mL/min for PAHs analysis and at 1.4 mL/min for
196 the other target compounds; and was held a constant flow by electronic pressure control. The
197 injector temperature was set to 300 °C. The split valve opened after 1 min, with a split ratio of
198 35/100. The manifold, ion trap source and transfer line temperatures were set to 120, 220 and
199 300 °C, respectively.

200 For the GC-MS analysis, except for organochlorines pesticides, all acquisition methods have
201 been previously described, the corresponding references being reported on Table 2 according
202 to the investigated chemical families. For HCB, γ -HCH, vinclozolin, metolachlor, α -
203 endosulfan, *o,p'*-DDT, β -endosulfan, methoxychlor and fenarimol analysis, the capillary
204 column was initially set at 80°C for thirty seconds, then ramped up to 280°C at 15 °C/min.
205 After 4.0 min at this value, the temperature was finally increased at 20 °C/min up to 350,
206 where it was held for 3.67 min. The total duration of GC analysis was 21 min. Ions were
207 formed under electro-ionization at 70 eV with an emission current of 80 μ A. The mass
208 spectrometer was operated using a hybrid acquisition mode alternating MS/MS and SIS
209 detection.

210 2.5.2. LC-MS/MS analyses

211 LC/MS is equipped with an electrospray ionization (ESI) source operated in negative-ion
212 mode. The analytical column used was a Pursuit C18 (150 mm x 2.1 mm I.D., 5µm particle
213 size, Varian, France) and a mobile phase consisting of acetonitrile (A) and water (B). The
214 gradient conditions were as followed: 25% of (A) and 75% of (B) for 2 min, followed by a
215 linear increase to 75% (A) at 16 min and 2 min hold at 75% (A). The flow rate was 0.2
216 mL/min and the injection volume was 20 µL. Only phytoestrogens were analysed by LC-MS.
217 The mass spectrometer was operated in order to record the MS/MS spectrum of deprotonated
218 pseudomolecular ion $[M-H]^-$. Collisions energies have been optimized between 20 to 30 eV
219 for each compound to obtain three characteristic ions. The optimized instrument conditions
220 were as followed: capillary voltage -3.4 kV, cone voltage, 50 V; multiplier voltage, 2250 V;
221 source temperature, 100°C; desolvation temperature, 450°C. Nitrogen was used as both the
222 nebulizing (50 L/h) and desolvation gas (700 L/h). Argon was used as collision gas at flow
223 0.28 (mL/min).

224

225 ***2.6. Data analysis and determination of bioassay- and instrumental-derived toxic-*** 226 ***equivalents***

227 Sigmoid dose-response curves and efficient concentrations (i.e. EC₂₅ and EC₅₀, corresponding
228 to concentrations of sediment extracts and chemical standards giving respectively 25 and 50%
229 of the maximum luciferase or EROD activities) were determined with the Regtox 7.5
230 Microsoft ExcelTM macro by using Hill equation (Vindimian et al., 1983) and freely available
231 at <http://eric.vindimian.9online.fr>. In all bioassays, significant responses were defined as those
232 greater than two times the standard deviation of the response obtained with DMSO (solvent
233 control). The dioxin-like, estrogenic, (anti)androgenic and PXR activities in samples derived
234 from bioassays were expressed as BaP- or TCDD-, E2-, DHT-, Flu- and SR12813-equivalents
235 (Bio-ref-EQs), respectively, which were determined as the ratio of the EC₂₅ of the reference

236 chemicals expressed as ng/L or µg/L to that of the sample expressed as equivalent gram of dry
237 weight sediment per litre (g EQ/L).

238 In order to estimate the contribution of analysed compounds to the total activity detected by a
239 bioassay, the measured concentrations (as ng/g sed wt) were converted to toxic-equivalent
240 activities derived from chemical analysis (Chem-ref-EQs), that is Chem-BaP-EQ, Chem-
241 TCDD-EQ, Chem-E2-EQ, Chem-DHT-EQ, Chem-Flu-EQ and Chem-SR12813-EQ, in each
242 respective bioassay. The Chem-ref-EQ were calculated according to the following equation:
243 $\text{Chem-ref-EQ} = \sum (C_i \times \text{ref-EF}_i)$, where, for a given chemical i , C_i is the measured
244 concentration in a sample and ref-EF_i is the inducing/inhibiting equivalent factor relative to
245 the reference ligand (i.e. BaP, TCDD, E2, DHT, Flu or SR12813) in a given bioassay. The
246 equivalent factors were calculated as follows: $\text{ref-EF}_i = \text{EC}_{25} \text{ of reference compound} / \text{EC}_{25} \text{ of}$
247 $\text{test compound } i$, on mass basis. Unless otherwise specified in Tables 3-5, ref-EFs of each
248 analysed compound were determined experimentally in each bioassay by establishing dose-
249 response curves for individual standard chemicals and by comparing them to that of the
250 reference compound. For instance, the 21 individual molecules listed in Table 4 were tested
251 for their ability to induce luciferase activity in MELN cells in our assay condition, and their
252 E2-equivalent factors (EEF) were then determined and used to calculate Chem-E2-EQ in the
253 samples. Finally, the ratio Chem-ref-EQ/Bio-ref-EQ allowed evaluating the contribution of
254 quantified compounds to the biological activity detected by the bioassay, i.e. a ratio near
255 100% means that all analysed compounds were explicative for biological results; otherwise
256 other non analysed compounds may be present in the samples.

257

258 **3. Results**

259 ***3.1. Dioxin-like activity in river sediment extracts***

260 The dose-response curves for EROD induction by all organic extracts are shown in Figure 1.
261 All samples elicited significant EROD activity, indicating a general contamination by dioxin-
262 like compounds in all studied sites. Sample responses varied greatly depending on the studied
263 site and exposure duration. Except for the Ais site after 24 h of exposure, the dose-response
264 curves were fairly complete and almost parallel to the dose-response curves for reference
265 chemicals. This allowed us to use EC₅₀ values for calculation of biological BaP-EQs and
266 TCCD-EQs (summarized in Table 3). The highest values were found for Lez site followed by
267 Rev, Rho, VdV and Ais. For all sediment extracts, EROD induction potency based on sample
268 concentration was higher after 4 h (Figure 1a) than after 24 h of exposure (Figure 1b) by
269 approximately one to three orders. This indicates a major contribution of non persistent
270 compounds, like PAHs, in the observed dioxin-like activities.

271 In order to test this hypothesis, the 16 priority PAHs were analysed in the extracts (Table 3).
272 Total PAHs concentrations indicate large between-site variations ranging from low
273 contamination levels in Ais site (0.22 µg/g d.w.) up to very high levels in Rev (11.98 µg/g
274 d.w.) and Lez (23.08 µg/g) d.w. sites. All sediment samples were dominated by high
275 molecular mass PAHs (4- to 6-ring).

276 The results given in Figure 2 showed that chemical- and bioassay-derived BaP-EQs and
277 TCCD-EQs were highly correlated ($r^2 = 0.99$ and 0.97 , respectively), which confirmed the
278 involvement of PAHs in the detected biological activities at both exposure durations.

279 However, in all sediment extracts, the Chem-EQs were significantly lower than the Bio-EQs,
280 indicating that only a part of activity was explained by analyzed PAHs, which accounted for
281 approximately 20 to 60 % and 16 to 40 % of the BaP-EQs and TCCD-EQs measured in
282 PLHC-1, respectively.

283

284 **3.2. Estrogenic activity in river sediment extracts**

285 The presence of estrogenic compounds in sediment extracts was tested in the MELN reporter
286 cell line (Figure 3). Significant dose-dependent induction of luciferase activity was obtained
287 for all tested sediment extracts; the response magnitudes varied between 34 % and 67 % of
288 the maximal response elicited by β -E2. Moreover, co-exposure with the pure anti-estrogen
289 ICI-182,780 led to inhibition of the luminescent signal (data not shown), thus showing the
290 specific involvement of the ER in the detected effects and indicating that estrogenic
291 compounds were present in samples. Because the slopes of dose-response curves for different
292 sediment extracts were not parallel to that of β -E2, the use of EC₂₅ was chosen to derive Bio-
293 E2-EQs. The Bio-E2-EQs values in the different extracts of sediment samples ranged from
294 0.20 to 6.43 ng/g d.w (Table 4), with the highest activity in the Réveillon sediment extract.
295 Overall, the results presented in Table 4 showed that E1, β -E2, 1OHPyr and BPA were
296 detected in all samples, while the synthetic estrogens were never detected. The Réveillon site
297 was the most contaminated by xeno-estrogens, with measurable levels of natural estrogens,
298 alkylphenols, PAH metabolites, parabens, chlorinated pesticides, as well as detectable levels
299 of the zearalenone metabolites α -ZAL and β -ZAL. α -E2 was detected at Rhonelle and Aisne
300 sites; equol was present at Rhonelle site, while other myco- and phyto-estrogens were not
301 detected.

302 In Aisne, Vallon du Vivier and Lézarde sediments, the Bio-E2-EQs were totally explained
303 only by the presence of E1 and β -E2. However, differences between chemical and biological
304 measurements were observed for Réveillon and Rhonelle sediments with Chem-E2-EQs/
305 Biol-E2-EQs ratio values of 28 and 35%, respectively. This suggests that these sediments may
306 contain estrogenic substances that were not included in our analytical methods.

307

308 ***3.3. (Anti) androgenic activity in river sediment extracts***

309 The dose-response curves show detectable androgenic activity in Aisne and Rhonelle samples
310 (Figure 4a), giving bioassay derived Bio-DHT-EQs values of 0.40 and 3.60 ng/g d.w.,
311 respectively. No androgenic activity was observed in the total sediment extract of Vallon du
312 Vivier, Lézarde and Réveillon sediments. In DHT co-exposure experiments (Figure 4b),
313 significant antiandrogenic activities were detected in Vallon du Vivier, Lézarde and Réveillon
314 samples, with respective Flu-EQs of 1.1, 7.4 and 32.5 µg/g d.w. By using the MTT test, no
315 cytotoxic effects could be observed for any working concentration of chemicals or organic
316 extracts (data not shown), thus indicating a specific effect on luciferase expression.
317 Determination of Chem-Flu-EQs showed that several anti-androgenic compounds were
318 present, but at concentrations that could explain only a very minor part of the observed
319 biological anti-androgenic activities (Table 5).

320

321 ***3.4. Detection of PXR ligands in river sediment extracts***

322 No anti-PXR activity was noted in these samples (data not shown). However, all five samples
323 induced PXR activation in a dose-response manner (Figure 5a). The response magnitudes
324 were always inferior to the maximum response produced by positive control (SR 12813 at 3
325 µM), and ranged from 30 % to 73 %. Nevertheless, luciferase induction was specific to PXR
326 activation since no luciferase increase was noted in the parental HG₅LN cell line, which
327 expresses only the GAL4-luciferase construct (Figure 5b). In HG₅LN cell line, inhibition of
328 luciferase at high concentrations of Rev and Lez extracts was noted and reflected early toxic
329 events.

330 The low Chem-SR12813-EQ/Bio-SR12813 ratios showed that 4tOP, 4nNP, βE2, E1, BPA,
331 endosulfan and *o,p'*-DDT, which are known PXR ligands, poorly contributed to Bio-
332 SR12813-EQs in the samples (Table 5). Thus, the detected PXR activities were due to other
333 non analysed compounds.

334

335

336 **4. Discussion**

337 *In vitro* profiling of sediment extracts showed multiple contaminations by dioxin-like and
338 endocrine active chemicals in small streams subjected to diffuse anthropogenic pollution. The
339 natural estrogens β E2 and E1 and of PAH-like compounds were identified as main
340 contributors to estrogenic and dioxin-like activities, respectively, as determined by the
341 bioassays. Conversely, (anti)androgenic and PXR-mediated activities were detected but the
342 responsible compounds could not be identified using targeted chemical analyses.

343 **4.1. PAHs and EROD inducing compounds**

344 To some extent, the levels of PAHs measured in this study were in line with previous studies
345 in French freshwater watersheds. For instance, total PAH content ranged from 1.2 to 12.5
346 $\mu\text{g/g}$ of dry sediment at various sites from the Seine estuary (Cachot et al., 2006), and from
347 2.3 to 41.3 $\mu\text{g/g}$ in the Seine and Marne rivers sampled at urban sites nearby Paris (Ollivon et
348 al., 2002). In our study, the highest PAH levels were detected in areas subjected to mixed
349 anthropogenic pressures like the Réveillon, located in urban area upstream of Paris, and the
350 Lézarde river, located nearby a highway road. Conversely, the Aisne site was located in a non
351 urbanized area and was logically found as low contaminated by PAHs.

352 In all samples, PAH-like compounds were major contributors of dioxin-like activity, which
353 mostly explained by PAHs with molecular weights (M.W.) of 228 and 252; however analyzed
354 PAHs explained only part of overall activity in PLHC-1 cells. This is not surprising since the
355 priority PAHs are used as qualitative tracers of pollution and we noted that other high
356 molecular weight PAHs, presumably active, were present in the samples (Figure 2). Indeed,
357 fullscan chromatogram of Lézarde extract (data not shown) showed the presence of detectable
358 concentrations of several isomers of Chry (M.W. 228) and BaP (M.W. 252) that probably

359 accounted for the detected activity in PLHC-1 cells, although this could not be tested in this
360 study. Similar conclusions have been reported by other studies that showed that a major
361 portion of AhR activity in river sediments was caused by nonpriority PAHs such as
362 methylated PAHs (e.g. Hollert et al., 2002; Brack et al., 2005). In addition, many other
363 dioxin-like compounds such as polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated
364 dibenzofurans (PCDFs), polychlorinated biphenyls (PCBs), polychlorinated naphthalenes
365 (PCNs) and some organochlorine pesticides, which were not included in the analyses, might
366 also explain the differences between chemical and biological analyses.

367 ***4.2. (Xeno) Estrogens***

368 The contamination of aquatic ecosystems by estrogenic compounds has been few reported in
369 France as compared to other industrialized countries. Nonetheless, effluents from sewage
370 water treatment plant (SWTP) have been shown to be the source of steroid estrogens at
371 different locations (Labadie and Budzinski, 2005, Muller et al., 2008; Cargouet et al., 2004);
372 in surface waters, natural and synthetic estrogens were also described as main estrogenic
373 chemicals in the Seine river downstream Paris, France (Cargouet et al., 2004). In sediments
374 sampled downstream from a large SWTP in the Seine river, Fenet et al. (2003) reported that
375 alkylphenols were the main contributors to the estrogenic activities detected by MELN
376 bioassay while steroid estrogens were not measured in their study. This contrasts with the
377 very low contribution of alkylphenols measured in the present study where estrogenic
378 activities were predominantly explained by the natural estrogens β E2 and E1. Since
379 alkylphenols are often considered as tracer compounds of water treatment effluents, the low
380 levels measured in our study could reflect the fact that our sites were not directly impacted by
381 SWTP effluents, the main source of aquatic pollution by alkylphenols. Nevertheless, other
382 studies in European countries have reported occurrence of steroid estrogens in river sediments
383 at similar levels than those found in our study, i.e. in the 0.3-5 ng/g range (Houtman et al.,

384 2006; Labadie and Hill, 2007; Reddy et al., 2005; Matějčiček et al., 2007). Moreover, the large
385 contribution of natural estrogens to biological E2-EQs is in agreement with the study of
386 Houtman et al. (2006) who reported that E1 and β E2 were responsible for more than 75 % of
387 the estrogenic activity in sediments from Zierikze harbour, The Netherlands. In the low
388 impacted sites (Aisne, Vallon du Vivier and Lézarde), natural estrogens nearly explained 100
389 % of the estrogenic activity detected by the bioassay. Conversely, in Réveillon and Rhonelle
390 samples, only a part (28 and 35 %) of the estrogenic activity was explained by β E2 and E1,
391 suggesting the presence of other xeno-estrogens in the samples. The Réveillon River is under
392 pressure of multiple sources of contamination and was indeed found to be contaminated by
393 many organic chemicals from agricultural (pesticides) and urban/industrial (BPA,
394 alkylphenols, parabens) origins. However, the targeted xeno-estrogens could not explain the
395 overall estrogenic of the sample, hence suggesting the presence of other ER ligands that were
396 not included in our analyses. This stresses the limits of the Chem-EQ based approach to
397 identify bioactive compounds in complex mixtures and argues for further investigations, for
398 instance by using effect directed analysis (EDA) approaches based on sample fractionation
399 and identification of bioassay active fractions (Brack, 2003), in order to better characterize the
400 contamination of this site by EDCs.

401 ***4.3. (Anti)androgenic activities***

402 The occurrence of (anti)androgenic compounds in river sediments has been rarely reported as
403 compared to estrogenic or dioxin-like compounds. However, such compounds can be present
404 in river sediment since high levels of androgenic activities (1-15 ng DHT-EQ/g) were
405 quantified by using the YAS assay in sediment from United Kingdom estuaries (Thomas et
406 al., 2002). Natural androgenic steroids in sewage treatment effluents were identified as
407 possible source of contamination (Thomas et al., 2002). Recently, Urbatzka et al. (2007)
408 reported both androgenic and antiandrogenic activities in fractionated sediment extracts from

409 the river Lambro, Italy, but the chemicals responsible for androgenicity in YAS assay were
410 not identified. Other studies have shown that pulp mill effluents (Jenkins et al., 2004) or
411 livestock feedlot effluents (Soto et al., 2004) are potential sources for androgens in the aquatic
412 environment. In our study, we detected androgenic activity (4 ng DHT-EQ/g) in Rhonelle
413 river sediment but the responsible compounds remain to be identified. In addition,
414 antiandrogenic activities were detected in three other sites (R ev, Lez and VdV). Although
415 several antiandrogenic chemicals were detected at these sites (i.e. BHT, BPA, alkylphenols,
416 pesticides, Table 5), the measured concentrations could explain only a minor part of the
417 activities measured in the MDA-kb2 bioassay. Therefore, the later were likely due to the
418 presence of other EDCs that have not been investigated in our chemical analyses.

419 ***4.4. PXR activities***

420 To our knowledge, the present study is the first demonstration of PXR-mediated activity in
421 river sediments. The human PXR is known to be activated by a large panel of environmental
422 chemicals that belong to different classes, like pharmaceuticals, steroids, alkylphenols (Mnif
423 et al., 2007), polybrominated diphenyl ethers (Pacyniak et al., 2007), as well as various
424 pesticides (Lemaire et al., 2006). In the present study, human PXR activating substances were
425 detected in all analyzed samples, at concentrations in the $\mu\text{g/g}$ range in terms of SR12813-
426 EQs. Since PXR shares several ligands with the estrogen receptor (ER), it was hypothesized
427 that several of the (xeno)estrogens detected in our samples could have contributed to the
428 PXR-mediated activity. However, the Chem-SR12813-EQ/Bio-SR12813-EQ ratios showed
429 that, at the measured concentrations, analyzed (xeno)estrogens only poorly contributed to the
430 SR12813-EQ quantities determined by the bioassay. Thus, more investigations using
431 fractionation and isolation procedures will be necessary to characterize the compounds
432 responsible for PXR activation in sediments. One promising methodology will likely consist
433 in the use of purified PXR immobilized on columns in order to isolate PXR ligands from

434 complex mixtures (Pillon et al., 2005; Balaguer et al., unpublished). It is expected that such
435 approach will allow identifying environmental PXR ligands and thus providing further useful
436 information on the toxicological relevance of detection of PXR activity in aquatic ecosystems.

437 ***4.5. Comparison with fish biomarkers***

438 In the present study, toxicological profiling of sediments showed the presence of a wide range
439 of chemicals that could potentially affect different molecular targets, namely ER, AR, AhR
440 and PXR, involved in the regulation of the endocrine system of exposed organisms. At the
441 studied sites, recent biomarker studies in wild three-spine stickleback (*Gasterosteus*
442 *aculeatus*) revealed fish exposure to different chemical stress including endocrine disrupters
443 and dioxin-like compounds (Sanchez et al., 2007, Sanchez et al., 2008). Although statistical
444 correlation between fish biomarker and *in vitro* bioassays could not be tested because of the
445 small number of sites, some concordances between the two approaches were noted. For
446 instance, vitellogenin induction in male stickleback has been evidenced in Réveillon and
447 Rhonelle rivers, which were the most active in the MELN assay (Table 4). In addition, female
448 stickleback from the Rhonelle River abnormally produced elevated levels of spigging, a male
449 glue protein synthesized in the kidney and used for building nest (Sanchez et al., 2008). This
450 suggested exposure to androgenic compounds, which correlates with the finding of AR-
451 mediated activity in MDA-kb2 cells in our study (Figure 4a). For dioxin-like activities,
452 significant EROD induction in male and female stickleback was reported in Réveillon and
453 Lézarde (Sanchez et al., 2007), the most active samples in the PLHC-1 assay (Table 3). On
454 the whole, the bioanalytical approach confirmed the multipollution context at impacted sites
455 like Réveillon and Rhonelle, and provided new information on possible causal agents for
456 abnormal endocrine responses in fish. Although we identified some of the chemicals
457 responsible for *in vitro* activities in sediment extracts, the direct extrapolation to fish exposure
458 is rather risky since it may depend on site specific characteristics that can influence pollutant

459 partitioning between sediment, suspended matter and dissolved phase in the water column.
460 Thus, more investigation using appropriate sampling methods like passive samplers will be
461 needed in order to link chemical contamination by EDCs and fish exposure and effects.

462 **5. Conclusion**

463 In summary, this study reports for the first time the simultaneous assessment of multiple
464 endocrine active and dioxin-like chemicals, in French river sediments sampled in small
465 streams subjected to various diffuse pollutions. Besides the major contribution of natural
466 steroids and PAH-like compounds to estrogenic and dioxin-like activities, (anti)androgenic
467 and PXR-mediated activities were detected although the individual active compounds could
468 not be identified using targeted chemical analyses. These samples are thus interesting
469 candidates for further EDA studies, which are under progress in order to elucidate the causal
470 agents.

471

472 **Acknowledgements**

473 The authors wish to thank Emmanuelle Maillot-Maréchal for excellent technical help with the
474 cell cultures. This study was funded by the French Ministry of Ecology and Sustainable
475 Development (Program 189), the “Agence Française de Sécurité Sanitaire de l’Environnement
476 et du Travail” (AFSSET, RD-2005-02) and by a doctoral fellowship from the ANRT and
477 INERIS (to SK).

478

479 **5. References**

480 Balaguer, P., Boussioux, A.M., Demirpence, E., Nicolas, J.C., 2001. Reporter cell lines are
481 useful tools for monitoring biological activity of nuclear receptor ligands. *Luminescence*
482 16, 153-158.

483 Brack, W., 2003. Effect-directed analysis: a promising tool for the identification of organic
484 toxicants in complex mixtures? *Analytical and Bioanalytical Chemistry* 377, 397-407.

485 Brack, W., Schirmer, K., Erdinger, L., Hollert, H., 2005. Effect-directed analysis of mutagens
486 and ethoxyresorufin-O-deethylase inducers in aquatic sediments. *Environmental*
487 *Toxicology and Chemistry* 24, 2445-2458.

488 Cachot, J., Geffard, O., Augagneur, S., Lacroix, S., Le Menach, K., Peluhet, L., Couteau, J.,
489 Denier, X., Devier, M.H., Pottier, D., Budzinski, H., 2006. Evidence of genotoxicity
490 related to high PAH content of sediments in the upper part of the Seine estuary
491 (Normandy, France). *Aquatic Toxicology* 79, 257-267.

492 Cargouet, M., Perdiz, D., Mouatassim-Souali, A., Tamisier-Karolak, S., Levi, Y., 2004.
493 Assessment of river contamination by estrogenic compounds in Paris area (France).
494 *Science of the Total Environment* 324, 55-66.

495 Eggen, R.I.L., Segner, H., 2003. The potential of mechanism-based bioanalytical tools in
496 ecotoxicological exposure and effect assessment. *Analytical and Bioanalytical Chemistry*,
497 377, 386-396.

498 Fenet, H., Gomez, E., Pillon, A., Rosain, D., Nicolas, J.C., Casellas, C., Balaguer, P., 2003.
499 Estrogenic activity in water and sediments of a French river: contribution of alkylphenols.
500 *Archives of Environmental Contamination and Toxicology* 44, 1-6.

501 Hartmann, N., Erbs, M., Wettstein, F.E., Schwarzenbach, R.P., Bucheli, T.D., 2007.
502 Quantification of estrogenic mycotoxins at the ng/L level in aqueous environmental
503 samples using deuterated internal standards. *Journal of Chromatography A*, 1138, 132-
504 140.

505 Hollert, H., Durr, M., Olsman, H., Halldin, K., Van Bavel, B., Brack, W., Tysklind, M.,
506 Engwall, M., Braunbeck, T., 2002. Biological and chemical determination of dioxin-like

507 compounds in sediments by means of a sediment triad approach in the catchment area of
508 the River Neckar. *Ecotoxicology* 11, 323-336.

509 Houtman, C.J., Cenijn, P.H., Hamers, T., Lamoree, M.H., Legler, J., Murk, A.J., Brouwer, A.,
510 2004. Toxicological profiling of sediments using *in vitro* bioassays, with emphasis on
511 endocrine disruption. *Environmental Toxicology and Chemistry* 23, 32-40.

512 Houtman, C.J., Booij, P., Jover, E., del Rio, D.P., Swart, K., van Velzen, M., Vreuls, R.,
513 Legler, J., Brouwer, A., Lamoree, M.H., 2006. Estrogenic and dioxin-like compounds in
514 sediment from Zierikzee harbour identified with CALUX assay-directed fractionation
515 combined with one and two dimensional gas chromatography analyses. *Chemosphere*, 65,
516 2244-2252.

517 Jenkins, R.L., Wilson, E.M., Angus, R.A., Howell, W.M., Kirk, M., Moore, R., Nance, M.,
518 Brown, A., 2004. Production of androgens by microbial transformation of progesterone *in*
519 *vitro*: A model for androgen production in rivers receiving paper mill effluent.
520 *Environmental Health Perspectives*, 112, 1508-1511.

521 Jobling, S., Williams, R., Johnson, A., Taylor, A., Gross-Sorokin, M., Nolan, M., Tyler, C.R.,
522 van Aerle, R., Santos, E., Brighty, G., 2006. Predicted exposures to steroid estrogens in
523 UK rivers correlate with widespread sexual disruption in wild fish populations.
524 *Environmental Health Perspectives* 114, 32-39.

525 Kavlock, R.J., Daston, G.P., DeRosa, C., Fenner-Crisp, P., Gray, L.E., Moore, J., Rolland, R.,
526 Scott, G., Sheehan, D.M., Sinks, T., 1996. A report of the US EPA-sponsored workshop.
527 *Environmental Health Perspectives* 107, 715-740.

528 Kinani, S., Bouchonnet, S., Bourcier, S., Creusot, N., Porcher, J.M., Ait-Aïssa, S., 2008a.
529 Extraction and purification procedures for simultaneous quantification of phenolic
530 xenoestrogens and steroid estrogens in river sediment by gas chromatography/ion trap
531 mass spectrometry. *Rapid Communications in Mass Spectrometry* 22, 3651-3661.

532 Kinani, S., Bouchonnet, S., Bourcier, S., Porcher, J.-M., Aït-Aïssa, S., 2008b. Study of the
533 chemical derivatization of zearalenone and its metabolites for gas chromatography-mass
534 spectrometry analysis of environmental samples. *Journal of Chromatography A* 1190,
535 307-315.

536 Labadie, P., Budzinski, H., 2005. Determination of steroidal hormone profiles along the Jalle
537 d'Eysines River (near Bordeaux, France). *Environmental Science & Technology* 39,
538 5113-5120.

539 Labadie, P., Hill, E.M., 2007. Analysis of estrogens in river sediments by liquid
540 chromatography-electrospray ionisation mass spectrometry - Comparison of tandem mass
541 spectrometry and time-of-flight mass spectrometry. *Journal of Chromatography A* 1141,
542 174-181

543 Laville, N., Aït-Aïssa, S., Gomez, E., Casellas, C., Porcher, J.M., 2004. Effects of human
544 pharmaceuticals on cytotoxicity, EROD activity and ROS production in fish hepatocytes.
545 *Toxicology* 196, 41-55.

546 Laville, N., Balaguer, P., Brion, F., Hinfrey, N., Casellas, C., Porcher, J.-M., Aït-Aïssa, S.,
547 2006. Modulation of aromatase activity and mRNA by various selected pesticides in the
548 human choriocarcinoma JEG-3 cell line. *Toxicology* 228, 98-108.

549 Lemaire, G., Mnif, W., Pascussi, J.M., Pillon, A., Rabenoelina, F., Fenet, H., Gomez, E.,
550 Casellas, C., Nicolas, J.C., Cavailles, V., Duchesne, M.J., Balaguer, P., 2006.
551 Identification of new human pregnane X receptor ligands among pesticides using a stable
552 reporter cell system. *Toxicological Sciences* 91, 501-509.

553 Louiz, I., Kinani, S., Gouze, M.E., Ben-Attia, M., Menif, D., Bouchonnet, S., Porcher, J.-M.,
554 Ben-Hassine, O.K., Aït-Aïssa, S., 2008. Monitoring of dioxin-like, estrogenic and anti-
555 androgenic activities in sediments of the Bizerta lagoon (Tunisia) by means of in vitro

556 cell-based bioassays: contribution of low concentrations of polynuclear aromatic
557 hydrocarbons (PAHs). *Science of the Total Environment* 402, 318-329.

558 Matejcek, D., Houserova, P., Kuban, V., 2007. Combined isolation and purification
559 procedures prior to the high-performance liquid chromatographic-ion-trap tandem mass
560 spectrometric determination of estrogens and their conjugates in river sediments. *Journal*
561 *of Chromatography A*, 1171, 80-89.

562 Mnif, W., Pascussi, J.-M., Pillon, A., Escande, A., Bartegi, A., Nicolas, J.-C., Cavallès, V.,
563 Duchesne, M.-J., Balaguer, P., 2007. Estrogens and antiestrogens activate hPXR.
564 *Toxicology Letters* 170, 19-29.

565 Mossman, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to
566 proliferation and cytotoxicity assays. *Journal of Immunology*, 65, 55-63.

567 Muller, M., Rabenoelina, F., Balaguer, P., Patureau, D., Lemenach, K., Budzinski, H.,
568 Barcelo, D., De Alda, M.L., Kuster, M., Delgenes, J.P., Hernandez-Raquet, G., 2008.
569 Chemical and biological analysis of endocrine-disrupting hormones and estrogenic
570 activity in an advanced sewage treatment plant. *Environmental Toxicology and*
571 *Chemistry* 27, 1649-1658.

572 Ohtake, F., Takeyama, K., Matsumoto, T., Kitagawa, H., Yamamoto, Y., Nohara, K.,
573 Tohyama, C., Krust, A., Mimura, J., Chambon, P., Yanagisawa, J., Fujii-Kuriyama, Y.,
574 Kato, S., 2003. Modulation of oestrogen receptor signalling by association with the
575 activated dioxin receptor. *Nature* 423, 545-550.

576 Ohtake, F., Baba, A., Takada, I., Okada, M., Iwasaki, K., Miki, H., Takahashi, S.,
577 Kouzmenko, A., Nohara, K., Chiba, T., Fujii-Kuriyama, Y., Kato, S., 2007. Dioxin
578 receptor is a ligand-dependent E3 ubiquitin ligase. *Nature* 446, 562-566.

579 Ollivon, D., Garban, B., Blanchard, M., Teil, M.J., Carru, A.M., Chesterikoff, C., Chevreuil,
580 M., 2002. Vertical distribution and fate of trace metals and persistent organic pollutants

581 in sediments of the Seine and Marne rivers (France). *Water Air and Soil Pollution* 134,
582 57-79.

583 Pacyniak, E.K., Cheng, X.G., Cunningham, M.L., Crofton, K., Klaassen, C.D., Guo, G.L.,
584 2007. The flame retardants, polybrominated diphenyl ethers, are pregnane X receptor
585 activators. *Toxicological Sciences* 97, 94-102.

586 Pillon, A., Boussioux, A.M., Escande, A., Aït-Aïssa, S., Gomez, E., Fenet, H., Ruff, M.,
587 Moras, D., Vignon, F., Duchesne, M.J., Casellas, C., Nicolas, J.C., Balaguer, P., 2005.
588 Binding of estrogenic compounds to recombinant estrogen receptor alpha : Application to
589 environmental analysis. *Environmental Health Perspectives*, 113, 278-284.

590 Reddy, S., Brownawell, B.J., 2005. Analysis of estrogens in sediment from a sewage-
591 impacted urban estuary using high-performance liquid chromatography/time-of-flight
592 mass spectrometry. *Environmental Toxicology and Chemistry* 24, 1041-1047.

593 Ryan, J.A., Hightower, L.E., 1994. Evaluation of heavy-metal ion toxicity in fish cells using a
594 combined stress protein and cytotoxicity assay. *Environmental Toxicology and*
595 *Chemistry* 13, 1231-1240.

596 Sanchez, W., Katsiadaki, I., Piccini, B., Ditche, J.M., Porcher, J.M., 2008. Biomarker
597 responses in wild three-spined stickleback (*Gasterosteus aculeatus* L.) as a useful tool for
598 freshwater biomonitoring: A multiparametric approach. *Environment International* 34,
599 490-498.

600 Sanchez, W., Aït-Aïssa, S., Palluel, O., Ditche, J.M., Porcher, J.-M., 2007. Preliminary
601 investigation of multi-biomarker responses in three-spined stickleback (*Gasterosteus*
602 *aculeatus* L.) sampled in contaminated streams. *Ecotoxicology* 16, 279-287.

603 Soto, A.M., Calabro, J.M., Prechtel, N.V., Yau, A.Y., Orlando, E.F., Daxenberger, A., Kolok,
604 A.S., Guillette, L.J., le Bizec, B., Lange, I.G., Sonnenschein, C., 2004. Androgenic and

605 estrogenic activity in water bodies receiving cattle feedlot effluent in eastern Nebraska,
606 USA. *Environmental Health Perspectives* 112, 346-352.

607 Sumpter, J.P., 2005. Endocrine disrupters in the aquatic environment: An overview. *Acta*
608 *Hydrochimica Et Hydrobiologica* 33, 9-16.

609 Thomas, K.V., Hurst, M.R., Matthiessen, P., McHugh, M., Smith, A., Waldock, M.J., 2002.
610 An assessment of in vitro androgenic activity and the identification of environmental
611 androgens in United Kingdom estuaries. *Environmental Toxicology and Chemistry* 21,
612 1456-1461.

613 Urbatzka, R., van Cauwenberge, A., Maggioni, S., Vigano, L., Mandich, A., Benfenati, E.,
614 Lutz, I., Kloas, W., 2007. Androgenic and antiandrogenic activities in water and sediment
615 samples from the river Lambro, Italy, detected by yeast androgen screen and chemical
616 analyses. *Chemosphere*, 67, 1080-1087.

617 Vethaak, A.D., Lahr, J., Schrap, S.M., Belfroid, A.C., Rijs, G.B.J., Gerritsen, A., de Boer, J.,
618 Bulder, A.S., Grinwis, G.C.M., Kuiper, R.V., Legler, J., Murk, T.A.J., Peijnenburg, W.,
619 Verhaar, H.J.M., de Voogt, P., 2005. An integrated assessment of estrogenic
620 contamination and biological effects in the aquatic environment of The Netherlands.
621 *Chemosphere* 59, 511-524.

622 Vindimian, E., Robault, C., Fillion, G., 1983. A method for co-operative and non co-operative
623 binding studies using non-linear regression analysis on a microcomputer. *Journal of*
624 *Applied Biochemistry*, 5, 261-268.

625 Vos, J.G., Dybing, E., Greim, H.A., Ladefoged, O., Lambre, C., Tarazona, J.V., Brandt, I.,
626 Vethaak, A.D., 2000. Health effects of endocrine-disrupting chemicals on wildlife, with
627 special reference to the European situation. *Critical Reviews in Toxicology* 30, 71-133.

628 Wilson, V.S., Bobseine, K., Lambright, C.R., Gray, L.E., 2002. A novel cell line, MDA-kb2,
629 that stably expresses an androgen- and glucocorticoid-responsive reporter for the

630 detection of hormone receptor agonists and antagonists. *Toxicological Sciences* 66, 69-
631 81.

632 **Figure legends**

633

634 Figure 1. Doses-response curves for EROD induction by sediment organic extracts in PLHC-1
635 cells measured after (a) 4h and (b) 24h exposure periods. Numbers correspond to the different
636 sites, with 1: Lez, 2: Rev, 3: Rho, 4: VdV, 5: Ais. Values represent the mean \pm SD, n=3.

637

638 Figure 2. Linear regression (log scale) showing correlation between PLHC-1 bioassay- (Bio-)
639 and PAH chemical analyses- (Chem-) derived BaP-EQs (a) and TCDD-EQs (b) in sediment
640 samples.

641

642 Figure 3. Estrogenic activity of (a) 17 β -estradiol and (b) the five sediment extracts (1: Rev, 2:
643 Rho, 3: VdV, 4: Ais, 5: Lez) in MELN cells. Results are expressed as percentage of maximal
644 luciferase activity induced by β -E2 at 10 nM. Values are means of triplicates \pm SD.

645

646 Figure 4. (a) Androgenic and (b) anti-androgenic activities of the reference chemicals
647 [dihydrotestosterone (DHT) and Flutamide (Flu)] and sediment extracts in MDA-kb2 cells.
648 Results are expressed as percentage of the maximal luciferase activity induced by DHT at (a)
649 10 nM and (b) 0.3 nM. Values are means of triplicates \pm SD.

650

651 Figure 5. (a) PXR-mediated dose-response curves of reference chemical (SR 12813) and
652 sediment extracts (1: Rev, 2: Rho, 3: Lez, 4: VdV, 5: Ais) in HG₅LN-PXR cells. Results are
653 expressed as percentage of maximal luciferase activity induced by SR 12813 at 0.3 μ M. (b)
654 Non specific effect of sediment extracts on constitutive luciferase expression in HG₅LN cells.
655 Results are expressed as percentage of luciferase activity in control cells. Values are means of
656 triplicates \pm SD.

657

658

659 Table 1. Summary of some general characteristics of sampling sites
 660

Sites	Aisne (Ais)	Vallon du Vivier (VdV)	Rhonelle (Rho)	Réveillon (Rev)	Lézarde Lez)
GPS coordinates	N 49°23'55" E 3°28'31"	N 49°43'23" E 0°27'42"	N 50°17'49" E 3°32'41"	N 48°34'00" E 2°32'09"	N 49°34'09" E 0°13'20"
Pressures ^a	Low	Mixed	Urban dense	Urban dense	Mixed
Water quality ^b	Good	Very good	Bad	Very bad	Good
Biomarker responses in fish (<i>Gasterosteus aculeatus</i>) ^c	No data	No alteration (reference site)	Induction of vitellogenin in male and spigging in female	Induction of EROD and vitellogenin in male	Induction of EROD

661
 662
 663 a, Data obtained from the Rivers Waterbase of the European Environmental Agency (EEA,
 664 2008); b, data from obtained from French water agencies; c, data from Sanchez et al. (2008).
 665

666 Table 2. Overview of investigated chemicals: chemical families, analytical standard sources
 667 and methods used for their quantification in samples
 668

<i>Classes (Providers)</i>	<i>Chemicals</i>	<i>Analytical methods (Reference)</i>
<i>Natural and synthetic estrogens</i> (Sigma-Aldrich)	estrone (E1), 17 β -estradiol (17β-E2), 17 α -estradiol (17α-E2), 17 α -ethynilestradiol (EE2), estriol (E3), mestranol (MeEE2), diethylstilbestrol (DES)	GC/MS with derivatization (Kinani et al., (2008a))
<i>Alkylphenols</i> (Sigma-Aldrich)	4-n-nonylphenol (4-n-NP), 4-tert-octylphenol (4-t-OP), 4-n-butoxyphenol (4-BuOP), 4,4'-isopropylidene diphenol (bisphenol A, BPA)	GC/MS/MS with derivatization (Kinani et al., (2008a))
<i>Parabens</i> (Sigma-Aldrich)	n-propylparaben (PrP), n-butylparaben (BP), benzylparaben (BzP)	GC/MS and GC/MS/MS with derivatization (Kinani et al., (2008a))
<i>hydroxy-PAHs</i> (Sigma-Aldrich)	1-hydroxypyrene (1-OHPyr), 2-hydroxyfluorene (2-OHFlu)	GC/MS/MS with derivatization (Kinani et al., (2008a))
<i>PAHs</i> (Sigma-Aldrich)	naphthalene (Nap), acenaphthylene (Acy), acenaphthene (Ace), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flt), pyrene (Pyr), benz[<i>a</i>]anthracene (B[a]A), chrysene (Chr), benzo[<i>b</i>]fluoranthene (B[b]F), benzo[<i>k</i>]fluoranthene (B[k]F), benzo[<i>a</i>]pyrene (B[a]P), indeno[1,2,3- <i>c,d</i>]pyrene (Ind), dibenz[<i>a,h</i>]anthracene (DBA), benzo[<i>g,h,i</i>]perylene (B[ghi]P)	GC/MS (Louiz et al., 2008)
<i>Organochlorine pesticides</i> (Promochem)	hexachlorobenzene (HCB), lindane (γ - HCH), vinclozolin, metolachlor, endosulfan (2 α :1 β), <i>o,p'</i> -DDT, methoxychlor, fenarimol	GC/MS and GC/MS/MS see section 2.5.1
<i>Phytoestrogens</i> (Sigma-Aldrich)	daidzein, genistein, biochanin A, equol, coumestrol, resveratrol	LC/MS/MS see section 2.5.2
<i>Mycoestrogens</i> (Sigma-Aldrich)	zearalenone (ZON), α -zearalenol (α-ZOL), β -zearalenol (β-ZOL), zearalanone (ZEA), α -zearalanol (α-ZAL) and β -zearalanol (β-ZAL)	GC/MS with derivatization (Kinani et al., (2008b))

669
 670
 671

672 Table 3. Concentrations of the 16 PAHs and their relative contribution to the total dioxin-like
 673 activity in sediment organic extracts from the five studied sites.
 674

Chemicals	LOQ ^a (ng/g)	IEF ^b		Concentration in sampling sites in ng/g d.w.				
		BaP 4h	TCDD 24h	Ais	VdV	Rho	Rev	Lez
Nap	0.11	n.i. ^c	n.i.	n.d. ^d	3.21	2.30	5.42	3.73
Acpy	0.17	5.56E-3	n.i.	n.d.	3.57	15.28	54.96	77.27
Acp	0.08	n.i.	n.i.	n.d.	4.94	2.56	26.18	38.66
Flu	0.09	1.44E-2	n.i.	n.d.	9.01	8.74	38.97	73.46
Phe	0.06	n.i.	n.i.	10.67	132.66	96.61	612.83	1706.67
Ant	0.11	n.i.	n.i.	3.30	37.44	46.01	225.27	712.69
Flt	0.05	n.i.	n.i.	46.35	307.42	267.01	1533.00	3367.66
Pyr	0.17	3.58E-3	3.85E-5	36.11	231.11	168.45	1307.67	2907.47
B[a]A	0.06	2.58E-1	9.77E-5	11.14	110.79	150.53	761.47	1477.33
Chr	0.06	2.92E-1	3.76E-4	8.02	102.24	172.16	1013.47	1596.00
B[b]F	0.13	6.94E-1	4.63E-4	15.59	69.66	723.05	2760.67	4674.00
B[k]F	0.11	2.94	4.23E-3	4.14	57.38	370.71	1602.09	2681.37
B[a]P	0.20	1	5.13E-4	15.26	97.99	138.03	841.60	1685.49
Ind	0.07	8.43E-1	1.64E-3	35.76	171.62	116.08	723.40	1190.89
DBA	0.12	3.66	6.11E-3	1.97	8.38	16.62	45.09	98.72
B[ghi]P	0.16	n.i.	n.i.	21.00	115.42	71.09	423.98	786.24
<i>Sum-PAHs (ng/g d.w.)</i>				<i>209.3</i>	<i>1462.8</i>	<i>236.5</i>	<i>11976.1</i>	<i>23077.7</i>
Chem-BaP-EQ (ng/g d.w.)				81	550	1978	8740	15037
Bio-BaP-EQ (ng/g d.w.)				200	910	7334	31556	75435
<i>Chem-BaP-EQ / Bio-BaP-EQ (%)</i>				<i>41</i>	<i>60</i>	<i>27</i>	<i>28</i>	<i>20</i>
Chem-TCDD-EQ (ng/g d.w.)				0.11	0.72	2.35	10.45	17.78
Bio-TCDD-EQ (ng/g d.w.)				0.67	4.49	5.89	38.43	48.38
<i>Chem-TCDD-EQ / Bio-TCDD-EQ (%)</i>				<i>16</i>	<i>16</i>	<i>40</i>	<i>27</i>	<i>37</i>

675
 676 ^a LOQ: limits of quantification provided by the GC-MS method, ^b IEF: induction equivalent factors relative to
 677 BaP after 4 h and to TCDD after 24 h of exposure (from Louiz et al., 2008), ^c n.i.: no EROD induction detected
 678 within the 0.1 nM to 10 µM concentration range, ^d n.d: not detected or below quantification limit.
 679

680 Table 4. Concentrations of estrogenic compounds measured in sediment organic extracts from
 681 the five studied sites, and their relative contribution to the total estrogenic activity measured
 682 in the MELN bioassay.
 683

Chemical classes	Chemicals	EEF ^a	LOQ ^b (ng/g)	Concentration in sediment extracts (ng/g d.w.)				
				Ais	VdV	Rho	Rev	Lez
<i>Natural estrogens</i>	α -E2	0.02	0.04	0.04	n.d.	0.2	n.d.	n.d.
	β -E2	1	0.05	0.27	0.78	0.58	1.58	0.18
	E1	0.02	0.02	0.3	0.58	0.48	1.28	0.36
	E3	0.17	0.6	n.d.	n.d.	n.d.	1.26	n.d.
<i>Synthetic estrogens</i>	EE2	0.93	0.07	n.d.	n.d.	n.d.	n.d.	n.d.
	MeEE2	0.02	0.08	n.d.	n.d.	n.d.	n.d.	n.d.
	DES	0.17	0.03	n.d.	n.d.	n.d.	n.d.	n.d.
<i>Alkylphenols</i>	4-t-OP	1.1E-04	0.03	n.d.	n.d.	0.56	6.3	n.d.
	4-n-NP	3.3E-06	0.01	n.d.	0.52	n.d.	0.04	n.d.
	BPA	4.5E-05	0.01	2.29	11.67	7.92	47.28	1.24
<i>Parabens</i>	PrP	7.4E-06	<0.01	n.d.	n.d.	n.d.	0.11	n.d.
	BuP	4.9E-06	<0.01	n.d.	n.d.	n.d.	0.07	n.d.
	BzP	6.6E-06	0.02	n.d.	n.d.	n.d.	0.79	n.d.
<i>PAHs metabolites</i>	1OHPyr	9.9E-07	0.02	0.11	0.26	0.3	0.41	0.37
	2OHFlu	6.3E-06	0.02	n.d.	n.d.	0.19	0.3	0.6
<i>Pesticides</i>	Endosulfan(α)	2.0E-06	0.21	n.d.	n.d.	n.d.	63.84	n.d.
	Endosulfan(β)	2.0E-06	1.25	n.d.	n.d.	n.d.	29.2	n.d.
	<i>o,p'</i> -DDT	1.7E-05	0.22	6.52	n.d.	n.d.	7.25	n.d.
<i>Phytoestrogen</i>	Equol	2.8E-04	0.15	n.d.	n.d.	0.17	n.d.	n.d.
<i>Mycoestrogens</i>	α -ZAL	0.14	0.13	n.d.	n.d.	n.d.	< LOQ	n.d.
	β -ZAL	0.03	0.02	n.d.	n.d.	n.d.	< LOQ	n.d.
Chem-E2-EQ (ng/g) ^d			-	0.28	0.79	0.59	1.82	0.19
Bio-E2-EQ (ng/g) ^e			0.010	0.29	0.83	1.69	6.43	0.20
Chem-E2-EQ / Biol-E2-EQ (%)				96	95	35	28	94

684
 685 ^a EEF: estradiol equivalence factors relative to estradiol, determined as described in the Materials and Methods
 686 section, ^b LOQ: limit of quantification, ^c n.d.: not detected or below quantification limits, ^d Chem-E2-EQs:
 687 chemical estradiol equivalents, ^e Biol-E2-EQs: biological estradiol equivalents based on EC₂₅ effective
 688 concentration in the MELN bioassay.
 689

690 Table 5. Summary of anti-androgenic and PXR-mediated activities in sediment extracts and
 691 relative contribution of known anti-androgenic and PXR ligands measured in organic extracts
 692 from the five studied sites.
 693

Chemicals	FEF ^a	SREF ^a	LOQ ^b (ng/g)	Concentration at sampling sites (ng/g sed d.w)				
				Ais	VdV	Rho	Rev	Lez
BPA	0.394	0.025	0.01	2.29	11.67	7.92	47.28	1.24
4-t-OP	0.339	0.099	0.03	n.d.	n.d.	0.56	6.3	n.d.
BHT	0.101	n.a.	0.07	n.d.	3.13	3.6	14.14	0.61
4-n-NP	0.016	n.a.	0.01	n.d.	0.52	n.d.	0.04	n.d.
Endosulfan(α)	0.058	0.029	0.21	n.d.	n.d.	n.d.	63.84	n.d.
<i>o,p'</i> -DDT	0.15	0.032	0.22	6.52	n.d.	n.d.	7.25	n.d.
Vinclozolin	2.68	n.a.	0.24	4.57	n.d.	n.d.	n.d.	n.d.
β -E2	n.a.	0.129	0.05	0.27	0.78	0.58	1.58	0.18
E1	n.a.	0.13	0.02	0.3	0.58	0.48	1.28	0.36
Chem-Flu-EQ (ng/g)				14.1	4.9	3.7	27.0	0.6
Bio-Flu-EQ (ng/g) ^d			17.0	nd	1089.4	nd	32493.4	7444.9
Chem-Flu-EQ / Biol-Flu-EQ (%)				-	0.45	-	0.08	0.01
Chem-SR12813-EQ (ng/g) ^d				0.34	0.47	0.39	4.26	0.10
Bio-SR12813-EQ (ng/g) ^d			40.0	964.8	1647.2	14322.2	51317.3	2147.1
Chem- SR12813-EQ / Biol- SR12813-EQ (%)				0.035	0.028	0.003	0.008	0.005

694
 695 ^a FEF (Flutamide equivalent factors) were determined as described in Materials and Methods; SREF
 696 (SR12813 equivalent factors) were from Lemaire et al. (2004) for endosulfan and from Mnif et al.
 697 (2007) for all other compounds. ^b LOQ: limit of quantification. n.a.: non active compound, n.d. not
 698 detected.
 699