

# 0-D-Modelling of Carbonaceous Aerosols over Greater Paris Focusing on the Organic Particle Formation

C. LIOUSSE<sup>1</sup>, C. MICHEL<sup>1</sup>, B. BESSAGNET<sup>2</sup>, H. CACHIER<sup>3</sup> and R. ROSSET<sup>1</sup>

<sup>1</sup>Laboratoire d'Aérodologie, CNRS-UPS, UMR 5560, 14 ave E. Belin, 31 400 Toulouse, France,  
e-mail: lioc@aero.obs-mip.fr

<sup>2</sup>INERIS, Direction des Risques Chroniques, Modélisation et Analyse Economique pour la gestion des Risques (MECO), Parc Technologique ALATA-BP n°2, 60550 Verneuil en Halatte, France

<sup>3</sup>Laboratoire des Sciences du Climat et de l'Environnement, CNRS-CEA, Ave de la Terrasse, 91 198-Gif sur Yvette Cedex, France

**Abstract.** To improve our understanding of aerosol formation and ageing in urban atmospheres, we have tested the ORISAM 0-D aerosol module (ORganic and Inorganic Spectral Aerosol Model). This module accounts for both types of primary carbonaceous particles (black carbon BC and primary organic carbon OC<sub>p</sub>) and also simulates the formation of secondary inorganic and organic particles (sulfates, nitrates, ammonium, water and secondary organic carbon particles OC<sub>sec</sub>) by attachment of gas precursors to pre-existing carbonaceous particles. Simulations were performed for surface aerosols over Greater Paris area during the ESQUIF summer 1998 and winter 2000 experiments. Results show that OC<sub>sec</sub> formation is highly dependent on temperature and insolation with more intense secondary formation in summer than in winter. Moreover in Summer, when atmospheric conditions shift from warm and humid to hot and dry, the model indicates a decreasing formation of secondary organic aerosols OC<sub>sec</sub> as shown by an increase of the OC<sub>p</sub>/(OC<sub>p</sub> + OC<sub>sec</sub>) ratio from 42 to 56%. These results satisfactorily compare with the few experimental available data for BC/(OC<sub>p</sub> + OC<sub>sec</sub>) ratios increasing from 24 to 37% against modelled values in the range 21–32%. ORISAM module sensitivity to initial size distributions, background concentrations and emissions of gases and primary carbonaceous particles was documented too. One main result is that the formation of secondary organic particles with ORISAM is very sensitive to the concentrations of gaseous precursors. At the present stage of ORISAM development, OC<sub>sec</sub> build up appears to be however less sensitive to particulate background concentrations.

**Key words:** carbonaceous aerosol, secondary organic aerosol formation, urban plumes, modelling

## 1. Introduction

Aerosol particles are often quoted for their overall cooling effect of about the same order of magnitude as the positive forcing by greenhouse gases (Charlson *et al.*, 1992; Haywood and Shine, 1995; Penner *et al.*, 1996; Jacobson, 2002; Chung and Seinfeld, 2002). However, these estimates are still subject to debate even in sign. The primary and most obvious reason is that the relative amount of the absorbing

fraction (mostly black carbon – BC) among scattering particles (sulphate, nitrate and organic particles) is poorly known. Also, more documentation is required, on the particle size distribution of each aerosol component, their mixing state (internal versus external) and subsequent optical properties. Moreover, due to the strong absorbing component of the urban aerosol mixture, top and bottom of the atmosphere radiative forcing values have been found quite different, involving important columnar heating rates (Léon *et al.*, 2003; Mallet, 2003).

Urban particles are expected to exert a significant direct forcing and to have a significant impact on the oxidising capacity of the atmosphere. They may also play a deleterious role on health (Atkinson *et al.*, 2001; Brauer *et al.*, 2003). Recently only, for urban atmospheres, the importance of carbonaceous particles and particularly of organic particles have been evidenced (Putaud *et al.*, 2003; Cachier *et al.*, 2005). This is partly due to important uncertainties founding the organic particle concentrations, uncertainties primarily related to specific artefacts encountered in both organic aerosol sampling and analysis procedures. Intercomparison of measurements (Schmidt *et al.*, 2001) and methods (<http://ocs.fortlewis.edu/Aerosols/OCEC/agenda.htm>) have shown that the determination of black carbon and organic carbon contents is operationally defined. Size distributions and the optical and hydrophilic properties of organic particles are also uncertain due to lack of data. Finally, emission inventories proposed in the literature for organic particles can be largely conflicting. A main reason relies in the primary or secondary origin of organic particles. Recent experiments have shown for example that the formation of secondary organic aerosol takes place readily close to sources (Cachier *et al.*, 2005). Due to this complexity, large discrepancies appear in the proposed global budgets if only the primary origin is considered (Cooke *et al.*, 1999), or the secondary origin (Kanakidou and Tsigaridis, 2000) or both (Lioussé *et al.*, 1996).

Aiming to better understand the formation of organic particles and to contribute to the construction of realistic budgets, a 0-D aerosol module ORISAM (ORganic and Inorganic Spectral Aerosol Model) was developed at Laboratoire d'Aérodologie (France). This aerosol module has first been used to simulate urban aerosol formation and transformation. Our urban case study applied to the Greater Paris area for summer 1998 and winter 2000 and relied on the ESQUIF experiment (Etude et Simulation de la Qualité de l'air en Ile de France (see <http://www.aero.jussieu.fr/themes/CA/Projet%20ESQUIF.html>)). The few existing measurements for carbonaceous aerosols were used to validate parameterizations and to run sensitivity tests.

## **2. The Modelling Procedure**

### **2.1. THE 0-D AEROSOL MODEL**

The ORISAM 0-D model is illustrated by the flow diagram in Figure 1. The module allows aerosol mixing and evolution by considering the formation of secondary




Figure 1. General schematic representation of ORISAM.

aerosols through coating of pre-existing insoluble aerosols. ORISAM relies on previous works of Bessagnet (2000) and Bessagnet and Rosset (2001). In our work, insoluble aerosols have been partitioned into black carbon (BC) and primary organic carbon (OC) and the secondary organic (OC<sub>sec</sub>) separated according to their origins (anthropogenic/biogenic origins). The secondary inorganic aerosols are sulphates, nitrates, ammonium and water. A sectional representation for aerosol size distribution is adopted with a variable number of sections (Fassi-Fihri *et al.*, 1997).

Coagulation (particle agglomeration), condensation, sedimentation and absorption of semi-volatile species are the different processes taken into account in the microphysical module. The RACM (Regional Atmospheric Chemistry Mechanism) scheme is used to describe the chemistry kinetics in the gas phase (Stockwell *et al.*, 1997). The chemical thermodynamic module ISORROPIA (Nenes *et al.*, 1998) is used to calculate for the inorganic components thermodynamic equilibrium between the particle and gas phases, most of them being highly dependent on atmospheric temperature and humidity. Secondary organic aerosols (SOA) are supposed to originate only from the absorption of volatile organic hydrocarbons onto the surface of primary particles. As it is not possible to account for all the complex, numerous and variable oxidation mechanisms taking place in the gas phase, some of them being even still unknown and in order to maintain reasonable computer time, SOA formation is parametrized. Following Wang *et al.* (1992), an alternative way is to simulate this formation by a two-step process: firstly, the total mass of condensable organic gases is obtained using an empirical method relying on experimental constant yields ( $Y$ ), and secondly, temperature-dependent partitioning coefficients ( $K_p$ ) (Odum *et al.*, 1996) are used to derive SOA mass.

## 2.2. AEROSOL MODEL INITIALIZATION

### 2.2.1. Period of Simulation

Two specific days were chosen during the summer 1998 dataset. Simulations were also conducted for 3 days of winter 2000 for sensitivity tests, though no experimental

data exist for this latter period. All selected days were chosen without rain and for their contrasted conditions (high relative humidity (RH) and low temperature, high temperature and low RH, low temperature and low RH).

### 2.2.2. Meteorological Data

Temperature, relative humidity, radiance, and boundary-layer thickness are necessary.

As shown in Figure 2 which display profiles of relative humidity and temperature, measured in Paris every hour and interpolated every minute for model input. August 7th, 1998, was quite hot (average temperature 23 °C) and dry (average relative humidity 47%). On the contrary, July 28th, 1998, was colder (17 °C), with reduced temperature diurnal amplitude, and much more humid (80%).

For winter 2000, January 22nd was cold (5 °C in the average) and humid (average relative humidity 81%); January 25th was still colder (−0.9 °C mean) and quite dry (49%), whereas January 30th was warmer (11 °C) and also quite humid (81%).

The boundary layer thickness ranged from 410 m to 1886 m in summer during ESQUIF, whereas 105–787 m for winter urban conditions. In the model, a sinusoidal function is used to reproduce the daily variation of the boundary layer thickness. The daily variation of radiances needed for the simulation, is inferred from the TUV radiation model (Tropospheric Ultraviolet and Visible radiation model), developed by Madronich and Flocke (1998).

### 2.2.3. Emissions

The same emissions are used for the different cases, whatever the season (summer and winter days) in order to focus on the impact of meteorological conditions only. The emissions of the anthropogenic gases (hydrocarbons, SO<sub>2</sub>, NH<sub>3</sub>, NO<sub>x</sub>), are extracted from the GENEMIS database (GENERation of an EMIssion database, <http://www.ier.unistuttgart.de/public/de/organisation/abt/tfu/projekte/genemis/GENEMIS.html>) with a 3 km spatial resolution. The gaseous species in the GENEMIS database differ from those introduced in the RACM regional model, so aggregation factors were used for adaptation (Stockwell *et al.*, 1997). The biogenic emissions (isoprene, monoterpenes) are obtained from a simulation for the Paris area performed with the Meso-NH-C model (Solmon, 2001). Let us note that, as described later in the paper, these choices are determinant to fix the background gaseous concentrations and then the SOA formation. Emission fluxes are given for 1-h time intervals for anthropogenic gases and 3-h time interval for biogenic gases, and subsequently interpolated to 1-min intervals.

BC and primary OC emission fluxes are obtained from a new inventory constructed for the year 1997 (Lioussé *et al.*, 2004). As no experimental data for carbonaceous aerosol size were available in ESQUIF we assumed the same distribution




Figure 2. Daily evolutions of temperature, relative humidity and  $OC_{sec}/OC_{tot}$  ratio (%) for (a) August 7th, 1998, (b) July 28th, 1998 and (c) January 25th, 2000.

as in Chazette and Liousse (2001). The uncertainty due to particle size is certainly rather high as shown by the preliminary sensitivity tests presented later in this paper. A two-mode log-normal distribution is considered. The first mode, assumed centred at  $0.1 \mu\text{m}$  (mass diameter size) contains only 50% BC mass. The second mode is centred at  $0.23 \mu\text{m}$  with 100% of primary OC mass and 50% of BC mass.

#### 2.2.4. Background Concentrations

Initial background gas concentrations were obtained at 00:00 local time (LT) after a spin-up period covering the whole day before. Different references for aerosol initialisation were chosen. Preliminary simulations presented in paragraph 3.1, point to high variabilities in BC, primary OC, secondary organic aerosol during the day and maximum concentrations at night (00:00 LT) due to accumulation processes. Indeed, no particle sink is taken into account in this modelling exercise (no particle deposition nor dynamical evolution except evolution of the boundary-layer thickness). Therefore, the mean state obtained by averaging aerosol concentrations the day before is that chosen for initialisation (called init-mean in the following). For example, primary carbonaceous aerosol concentrations for the 08/07/98 are  $1.53 \mu\text{g}/\text{m}^3$  (50% BC) for the first mode and  $7.87 \mu\text{g}/\text{m}^3$  (50% BC and 100% OC) for the second mode.

### 3. Modelling Results


#### 3.1. SIMULATED AEROSOL EVOLUTION

Figures 3a–c display the evolution of the different aerosol components obtained respectively for 3 days: August 7th and July 28th, 1998 and January 25th, 2000. Results are presented at 0:00, 12:00 and 24:00 LT. The secondary aerosols either inorganic (sulphates, nitrates, ammonium and water) or organic (SOA) display increasing concentrations with the hour of the day, and marked differences from one meteorological situation to another.


- Concentrations of the secondary inorganic aerosols are quite different for winter and summer days, and appear to be mainly driven by the competitive effects of relative humidity, temperature and photochemical conditions. As shown with the results for summer days (Figures 3a and b), formation of the inorganic aerosol components is twice more important on 07/28/98 ( $28 \mu\text{g}/\text{m}^3$ ) as compared to 08/07/98 ( $13 \mu\text{g}/\text{m}^3$ ), when afternoon relative humidity is higher and temperature lower. The water capture may cause enhancement of the absorption of the different semi-volatile species at the aerosol surface, and thus the formation of secondary inorganic species. The role of photochemistry is illustrated through comparison between the two days January 25th and August 7th with about the same mean atmospheric relative humidity (47 and 49%): a significant difference

**Mass size distribution**


**3a : 08/07/98**



**3b : 07/28/98**



**3c : 01/25/00**



*Figure 3.* Evolution of the different aerosol components respectively for (a) August 7th, 1998, (b) July 28th, 1998 and (c) January 25th, 2000 (P1: black carbon, P2: Primary organic carbon, M: water soluble fraction including water, Sec-ant-aero: Anthropogenic Secondary organic aerosol). Sec-bio-aero: Biogenic Secondary organic aerosol).

is there observed for the inorganic aerosol components ( $7.5$  and  $13 \mu\text{g}/\text{m}^3$ ) which points to the importance of photochemistry during summer days. This sensitivity to photochemical formation also exists at higher relative humidities (RH about 80%), similar differences being observed in winter ( $11$ – $12.6 \mu\text{g}/\text{m}^3$  for

simulations performed on January 22nd and 30th (results not shown here)) and in summer ( $28 \mu\text{g}/\text{m}^3$  for simulations of July 28th).

- The formation of secondary organic aerosol generally begins in the early morning due to the conjunction of the morning traffic peak emissions and the onset of photochemistry. When temperature reaches a maximum, secondary aerosol VOC precursors are found predominantly in the gas phase. When the temperature decreases, around 18:00 LT approximately, the transfer of condensable organic species from the gas to the particulate phase may become more important. Such a process, driven by the module parameterizations, is in agreement with the theoretical works of Sheehan and Bowman (2001), who have shown that temperature governs the transfer of organic species from the gas to the particulate phase and conversely. This temperature dependence is illustrated by comparison of the two summer days in which the mean temperature is markedly different: when comparing Figures 2a and b to Figure 3a and b, it may be seen that, with almost similar afternoon photochemistry activity for the two summer days, the relative concentration of secondary organic carbon is weaker on August 7th, 1998 than for the milder day, July 28th 1998. Such a difference is still enhanced after 24 h of simulation ( $11.4$  and  $7.65 \mu\text{g}/\text{m}^3$  on July 28th and August 7th). During the hot and dry day August 7th, SOA formation is reduced when temperature reaches a maximum, as VOCs remain in the gaseous phase. At the end of the afternoon, when temperature decreases, SOA formation again increases, a process which may be favoured by the parallel increase of traffic precursor emission. This SOA formation is still more visible due to the decrease of the boundary layer thickness. For the selected winter days, similar profiles of secondary organic aerosol formation are obtained. However, SOA concentrations are weaker by a factor of two (mean:  $4.6 \mu\text{g}/\text{m}^3$  in winter whereas  $9.5 \mu\text{g}/\text{m}^3$  in summer), which may primarily reflect the impact of photochemistry.

### 3.2. COMPARISON BETWEEN EXPERIMENTAL AND MODELLED VALUES

Table I presents  $\text{BC}/\text{OC}_{\text{tot}}$  ratio values simulated for the different situations, in winter and summer. It may be recalled that  $\text{OC}_{\text{tot}}$  includes both  $\text{OC}_{\text{p}}$  and  $\text{OC}_{\text{sec}}$ . Measured values are shown for the summer experiment. As BC is an inert species, and considering that emissions are the same for the different case studies, any decrease of  $\text{BC}/\text{OC}_{\text{tot}}$  ratio may be interpreted as an indicator of secondary OC production.

A statistically significant difference is observed for mean  $\text{BC}/\text{OC}_{\text{tot}}$  ratios obtained by the model in winter (43%) and in summer (33%). In winter, lower temperatures and weak radiances create conditions of low photolytic activity of and consequently, weaker SOA formation than in summer. Higher  $\text{BC}/\text{OC}_{\text{tot}}$  ratios are thus expected in winter. Interestingly such a seasonal trend in  $\text{BC}/\text{OC}_{\text{tot}}$  ratios was experimentally obtained by Castro *et al.* (1999) for urban areas in Portugal in summer 1993 and winter 1992–1993 in Coimbra (33 and 48% respectively) and in Oporto (37 and 59%).



Table I. Simulated BC/OC<sub>tot</sub> ratios (%) for the selected winter and summer days

Date	Simulated BC/OC <sub>tot</sub> ratio (%)				Measured BC/OC <sub>tot</sub> ratio (%) 9–21 h
	Mean	0–9 h	9–21 h	21–24 h	
01/22/00	45	49	43	42	No data
01/25/00	45	48	42	42	No data
01/30/00	40	45	37	36	No data
08/07/98	37	42	<b>37</b>	21	<b>32</b>
07/28/98	29	40	<b>24</b>	20	<b>21</b>

Note. Summer days with existing experimental data are in bold.

An interesting feature in Table I is also that modelled average values (“mean day values” for the 9–21 h LT time period) compare satisfactorily with experimental values obtained during the summer ESQUIF campaign especially for their trends. Indeed, BC/OC<sub>tot</sub> ratio values for July 28th and August 7th show the same decrease of a third for either measurements or simulated values. If we recall the important discrepancies appearing for urban carbonaceous aerosol measurements between different analytical protocols (Schmidt *et al.*, 2001), this fair agreement between simulated and experimental values may be coincidental but similar trends of BC/OC<sub>tot</sub> under different meteorological situations is certainly an indicator of a reasonable parameterisation of the SOA formation in ORISAM for our case study. Furthermore, the experimental values were obtained for Gif sur Yvette aerosols at a suburban location, whereas the modelled values were calculated for central Paris. This difference in location has been shown to introduce a small shift of about 3% in experimental BC/OC ratios (Ruellan and Cachier, 2001), the pure urban aerosol being enriched in BC. This minor correction still emphasizes the agreement. However, due to the quoted limitations (linked to both experiments and modelling) more data comparisons are needed to actually validate the model SOA formation.

### 3.3. SENSITIVITY TESTS UPON INITIALISATION CONDITIONS

#### 3.3.1. Impact of Temperature and Relative Humidity on SOA Formation

Two sets of sensitivity tests have been conducted with the same emissions and initialisations, changing the temperature or the relative humidity only. Figure 4 displays 24 h averages of values of OC<sub>sec</sub>/OC<sub>tot</sub> ratios obtained for the different simulations. Although comparison is based on 24 h averages, OC<sub>sec</sub> formation is highly dependent on temperature. However, dependency with the relative humidity is seen to be less important.

Interestingly, a temperature threshold seems to appear around 20 °C, with a sharp decrease of the OC<sub>sec</sub>/OC<sub>tot</sub> ratios around this temperature value. At lower




Figure 4. Temperature threshold for SOA formation as seen by the evolution of the OC<sub>sec</sub>/OC<sub>tot</sub> ratio, for different meteorological conditions: two relative humidity (50 and 80%) and two seasonal (winter and summer) insolation conditions.

temperatures, increasing temperature favours the SOA formation with although much less effect. This threshold is apparently the same for both seasons but as expected, photochemical OC<sub>sec</sub> formation is more significant in summer than in winter (up to 50% more). Such results confirm the hierarchy of processes described in Section 3.1.

### 3.3.2. Sensitivity Tests for SOA Formation According to Aerosol and Gaseous PreCursor Initialization

Different tests were conducted to capture the role of initial concentration levels of aerosol and gaseous precursors. Firstly, the role of the carbonaceous particle size distribution was scrutinized. This parameter is probably the one subject to the larger uncertainties, due to lack of measurements. For the purpose of the tests, particle sizes used for “actual conditions” (Section 2.2: 0.1  $\mu\text{m}$  for the first mode and 0.23  $\mu\text{m}$  for the second mode) were respectively divided or multiplied by a factor of two. Such a range may account for more recent results (Putaud *et al.*, 2003; Cachier *et al.*, 2005). It was found in our sensitivity tests however, that these changes of particle size have no impact on OC<sub>sec</sub> formation whereas important changes of the particle specific surface are expected with changes in the particulate size distribution.

Secondly, as primary aerosol and gas concentrations have probably a strong impact on SOA formation, different simulations were performed changing concentrations of either the carbonaceous particles or the gases. Figures 5a and b show the relative increase in secondary organic aerosol due to changes in emissions when aerosol (Figure 5a) and gas amounts (Figure 5b) are successively divided and




Figure 5. Relative increase of OC<sub>sec</sub> (%). (a) For different particle emissions and (b) for different gas emissions.

multiplied by a factor of 8, 4 or 2. As expected, it is shown that the secondary aerosol formation is highly dependent on the initial concentrations of aerosols or precursor gases. Also, it may be seen that the response to gas changes is much more important than the one obtained through changes in aerosol concentration levels: the range varies from -85% to almost 300% for gases whereas only from -14% to 33% for particles. It is interesting to note that significant differences occur between the two meteorological situations illustrated in Figure 5a. On 07/28/98, increase or decrease in particle emissions have a weaker impact on SOA formation when compared to 08/07/98. The impact of aerosol concentrations thus seems to be more important for the day with less favourable conditions (high temperature, low relative humidity) for SOA formation. Indeed the number of aerosols might play an

Table II. Simulated BC/OC<sub>tot</sub> ratios (%) for two different initial background concentration conditions at 0:00

Initialisation type	Date (9–21 h)	Mean simulated BC/OC <sub>tot</sub> ratio (%)	Mean measured BC/OC <sub>tot</sub> ratio (%)
Init-mean	01/25/00	42	No data
	08/07/98	37	32
	07/28/98	24	21
Init-norm	01/25/00	46	No data
	08/07/98	23	32
	07/28/98	20	21

Note. “Init-mean” with the average daily concentrations obtained by a 24-h spin-up the day before. “Init-norm” with concentrations obtained at 00:00 after a 24-h spin-up the day before.

important role to allow the transfer of the high gaseous organic precursors amount into particulate phase. The situation is different in Figure 5b where conditions of SOA formation remain more favourable on 07/28/98 than on 08/07/98 (more hot, less humid), whatever the initialisation.

The initialisation procedure (called init-mean) relies on average background aerosol concentration values obtained by a 24 h spin-up performed for the previous day. Sensitivity tests were conducted by changing background concentrations of primary BC and OC particles by applying the init-norm test. Let us recall that the init-norm test refers to an initialisation with instantaneous concentrations taken at 24:00 LT at the end of the spin-up. Comparisons of BC/OC<sub>tot</sub> ratios obtained for the two different initialisations are shown in Table II. The initialisation procedure may be seen to have an important impact on BC/OC<sub>tot</sub> values.

For summer, BC/OC<sub>tot</sub> ratios are much lower in the new (init-norm case) simulation than in the previous (init-mean case) one. This is mainly due to higher primary organic aerosol concentrations for the new case: BC concentrations are multiplied by 1.5 (for August 7th and July 28th), OC<sub>p</sub> by 2.1 and 2.5 (for August 7th and July 28th, respectively). In addition, differences in OC<sub>sec</sub> concentrations between the two cases are small as compared to differences in primary organic aerosol concentrations. Indeed, differences between secondary aerosol formation rates between init-norm and init-mean situations are slightly negative (−15 and −37%), which means that the secondary aerosol formation is favoured under the mean (not extreme) conditions. The difference between the two initialisation conditions is less pronounced for 07/28/98 than for 08/07/98. It seems that for this day (07/28/98) with particularly optimal conditions for secondary organic carbon formation, the initialisation conditions are less influent.

For winter days, the variation in primary aerosol concentrations between the two initialisations is of the same order, 1.4 both for BC and OC<sub>p</sub>. Same tendency is observed for the two other winter days.

Finally, it was observed that comparisons of simulations with experimental data are worst for all days with the init-norm conditions.

#### 4. Conclusion

Using the 0-D ORISAM aerosol module, tests were performed to simulate aerosol formation and ageing over Paris during the ESQUIF experiment for summer 1998 and winter 2000.

The module parameterizations allow to reproduce trends in the formation of secondary organic particles (SOA) and differences found experimentally between summer and winter aerosols. Moreover, in spite of the difficulty to accurately obtain actual BC and OC aerosol contents (still subject to debate), the few experimental black carbon to organic carbon ratio values available for this comparison are well reproduced for the summer season. No validation has been possible for winter due to lack of measurements, though there is still a general agreement for qualitative comparisons.

The module sensitivity to different changes was tested, such as temperature, relative humidity, initial size distribution, background particulate concentrations and emissions of gases and primary carbonaceous aerosol. The most important effect is certainly that of temperature. This temperature effect shows a marked threshold value (20 °C) at both seasons above which VOC transfer from the gas to particulate phase could be reduced, thus precluding any further important SOA formation. Further tests are ongoing to check if this threshold value is found at other sites. Changes in gas emissions have been shown to have a large impact on SOA formation too. Further investigations are needed to test the relative importance of chemically different VOC precursors. Unexpectedly, the aerosol size distribution has no quantifiable impact on the SOA formation, so further developments should be proposed for the parameterisation of OC<sub>sec</sub> formation to account for the role of particle surface areas, in particular effective thermodynamic links of inorganic and organic species and particles (Pun *et al.*, 2002).

Changes in aerosol number appear of poor importance and a weak effect only of particulate background concentrations may be observed. It may be hypothesized that under unfavourable hot conditions when the gas to particle transfer is slowed down by high temperature, the need of aerosol sites active for absorption processes, could be then more determinant.

This preliminary work needs to be confirmed and completed by other studies over different areas in order to conclude if this type of module is able to reproduce SOA formation in a wide range of situations without considering more complex parameterisations. Tests on the impact of different combustion sources (biogenic/anthropogenic, fossil fuel/biomass burning) are ongoing. Furthermore, 1D and 3D aerosol modelling studies with ORISAM have allowed to test successfully along with the particulate phase transformations, the particle transport as well as wet and dry depositions (Cousin *et al.*, 2005).

## References

- Atkinson, R. W., Anderson, H. R., Sunyer, J., Ayres, J., Baccini, M., Vonk, J. M., Boumghar, A., Forastiere, F., Forsberg, B., Touloumi, G., Schwartz, J. and Katsouyanni, K., 2001: Acute effects of particulate air pollution on respiratory admissions: Results from the APHEA 2 project. *Air pollution and health: A European approach, Am. J. Respir. Crit. Care Med.* **164**(10 Pt 1), 1860–1866.
- Bessagnet, B., 2000: Modélisation de l'aérosol carboné – cas particulier des particules fractales de combustion. PhD Thesis. Université Paul Sabatier. Toulouse.
- Bessagnet, B. and Rosset, R., 2001: Fractal modelling of carbonaceous aerosols-application to car exhaust plumes, *Atmos. Environ.* **35**, 4751–4762.
- Brauer, M., Hoek, G., van Vliet, P., Meliefste, K., Fischer, P., Gehring, U., Heinrich, J., Cyris, J., Bellander, T., Lewne, M., and Brunekreef, B., 2003: Estimating long-term average particulate air pollution concentrations: Application of traffic indicators and geographic information systems, *Epidemiology* **14**(2), 228–239.
- Cachier, H., Aulagnier, F., Sarda, R., Gautier, F., Masclet, P., Besombes, J.-L., Marchand, N., Despiiau, S., Croci, D., Mallet, M., Laj, P., Marinoni, A., Deveau, P.-A., Roger, J.-C., Putaud, J.-P., Van Dingenen, R., Dell'Acqua, A., Viidanoja, J., Martins-Dos Santos, S., Liousse, C., Cousin, F., and Rosset, R., 2005: Aerosol studies during the ESCOMPTE experiment: An overview, *Atmos. Res.* **74**, 547–563.
- Castro, L. M., Pio, C. A., Harrison, R. M., and Smith, S. J. T., 1999: Carbonaceous aerosol in urban and rural european atmospheres. Estimation of the secondary organic carbon concentrations, *Atmos. Environ.* **31**, 2771–2781.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Jr, Hansen, J. E., and Hofmann, D. J., 1992: Climate forcing by anthropogenic aerosols, *Science* **255**, 423–430.
- Chazette, P. and Liousse, C., 2001: A case study of optical and chemical ground apportionment for urban aerosols in Thessaloniki, *Atmos. Environ.* **35**, 2497–2506.
- Chung, S. H. and Seinfeld, J. H., 2002: Global distribution and climate forcing of carbonaceous aerosols, *J. Geophys. Res.* **107**(D19), 4407.
- Cooke, W. F., Liousse, C., and Cachier, H., 1999: Construction of a  $1^\circ \times 1^\circ$  fossil fuel emission data set for carbonaceous aerosol and implementation and radiative impact in the ECHAM4 model, *J. Geophys. Res.* **104**(22), 137–162.
- Cousin, F., Liousse, C., Cachier, H., Bessagnet, B., Guillaume, B., and Rosset, R., 2005: Aerosol modeling and validation during ESCOMPTE 2001, *Atmos. Environ.* **39**, 1539–1550.
- Fassi Fihri, A., Suhre, K., and Rosset, R., 1997: Internal and external mixing in atmospheric aerosols by coagulation: Impact on the optical and hygroscopic properties of the sulphate-soot system, *Atmos. Environ.* **31**, 1393–1402.
- Haywood, J. M. and Shine, K. P., 1995: The effect of anthropogenic sulfate and soot aerosol on the clear sky planetary radiation budget, *Geophys. Res. Lett.* **22**(5), 603–606.
- Jacobson, M. Z., 2002: Control of fossil-fuel particulate black carbon and organic matter, possibly the most effective method of slowing global warming, *J. Geophys. Res.* **107**, 4410.
- Kanakidou, M. and Tsigaridis, K., 2000: Human activity enhances the formation of organic aerosols by biogenic hydrocarbon oxidation, *J. Geophys. Res.* **12**, 9243–9254.
- Léon, J. F., Chazette, P., Pelon, J., Dulac, F., and Randriamarisoa, H., in press: Aerosol direct radiative forcing impact over the INDOEX area based on active and passive remote sensing, *J. Geophys. Res.*
- Liousse, C., Guillaume, B., Michel, C., Junker, C., Grégoire, J. M., and Cachier, H., 2004: *Recent Developments for Carbonaceous Aerosol Inventories of the 1950–2100 Time Period*, International Carbonaceous Aerosol Conference, Vienna.

- Lioussé, C., Penner, J. E., Chuang, C., Walton, J. J., Edleman, H., and Cachier, H., 1996: A global three-dimensional model study of carbonaceous aerosols. *J. Geophys. Res.* **101**, 19411–19432.
- Madronich, S. and Flocke, S., 1998: The role of solar radiation in atmospheric chemistry, in P. Boule (ed.), *Handbook of Environmental Chemistry*, Heidelberg, Springer-Verlag, pp. 1–26.
- Mallet, M., 2003: Etude des propriétés microphysiques et optiques de l'aérosol en zone péri-urbaine durant Escompte. Estimation de l'impact radiatif direct. PhD thesis, University of Toulon and Var, 135 pp.
- Nenes, A., Pilinis, C., and Pandis, S. N., 1998: ISORROPIA: A new thermodynamic model for inorganic multicomponent atmospheric aerosols, *Aquat. Geochem.* **4**, 123–152.
- Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H., 1996: Gas/particle partitioning and secondary aerosol yield, *Environ. Sci. Technol.* **31**, 1890–1897.
- Penner, J. E., Chuang, C. C., and Lioussé, C., 1996: The contribution of carbonaceous aerosols to climate change, in M. Kulmala and P. E. Wagner (eds.), *Nucleation and Atmospheric Aerosols*, Elsevier, New York, pp. 759–769.
- Pun, B. K., Griffin, R. G., Seigneur, C., and Seinfeld, J. H., 2002: Secondary organic aerosol: 2. Thermodynamic model for gas/particle partitioning of molecular constituents, *J. Geophys. Res.* **107**, 4333, doi:10.1029/2001JD000542.
- Putaud, J.-P., Baltensperger, U., Brüggemann, E., Facchini, M. C., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Jones, A. M., Laj, P., Maenhaut, W., Mihalopoulos, N., Müller, K., Palmgren, F., Querol, X., Rodriguez, S., Spindler, G., ten Brink, H., Tunved, P., Van Dingenen, R., Wehner, B., Weingartner, E., Wiedensohler, A., Wahlin, P., and Raes, F., 2003: A European aerosol phenomenology. Physical and chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, Joint Research Centre, European Commission, EUR 20411 EN.
- Ruellan, S. and Cachier, H., 2001: Characterisation of fresh particulate vehicular exhausts near a Paris high flow road, *Atmos. Environ.* **35**, 453–468.
- Schmidt, H., Laskus, L., Abraham, H. J., Baltensperger, U., Lavanchy, V., Bizjak, M., Burba, P., Cachier, H., Crow, D., Chow, J., Gnauk, T., Even, A., Brink, H. M. T., Giesen, K. P., Hitzinger, R., Hueglin, C., Maenhaut, W., Pio, C., Carvalho, A., Putaud, J. P., Toom-Sauntry, D., and Puxbaum, H., 2001: Results of the “carbon conference” international aerosol carbon round robin test stage 1, *Atmos. Environ.* **25**, 2111–2121.
- Sheehan, P. E. and Bowman, F. M., 2001: Estimated effects of the temperature on secondary organic aerosol concentrations, *Environ. Sci. Technol.* **35**(11), 2129–2135.
- Solmon, F., 2001: Modélisation méso-échelle des émissions biogéniques d'hydrocarbures volatiles en France. Thesis. Institut Nationale d'Agronomie. Paris-Grignon.
- Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S., 1997: A new mechanism for regional atmospheric chemistry modeling, *J. Geophys. Res.* **102**, 25847–25879.
- Wang, S. C., Paulson, S. E., Grosjean, D., Flagan, R. C., and Seinfeld, J. H., 1992: Aerosol formation and growth in atmospheric organic/NO<sub>x</sub> systems – I. Outdoor smog chamber studies of C<sub>7</sub>- and C<sub>8</sub>-hydrocarbons, *Atmos. Environ. A* **26**, 403–420.