



Multi-model assessment of PM trends in europe during two decades (1990-2010)

Svetlana Tsyro, Camilla Andersson, Bertrand Bessagnet, Augustin Colette, Florian Couvidat, Kees Cuvelier, Astrid Manders, Kathleen Mar, Mihaela Mircea, Noelia Otero, et al.

► To cite this version:

Svetlana Tsyro, Camilla Andersson, Bertrand Bessagnet, Augustin Colette, Florian Couvidat, et al.. Multi-model assessment of PM trends in europe during two decades (1990-2010). 18. International Conference on Harmonisation within Atmospheric Dispersion Modelling for Regulatory Purposes (HARMO 18), Oct 2017, Bologne, Italy. pp.61-64. ineris-01863207

HAL Id: ineris-01863207

<https://hal-ineris.archives-ouvertes.fr/ineris-01863207>

Submitted on 28 Aug 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTI-MODEL ASSESSMENT OF PM TRENDS IN EUROPE DURING TWO DECADES (1990-2010)

Svetlana Tsyro¹, Camilla Andersson², Bertrand Bessagnet³, Augustin Colette³, Florian Couvidat³, Kees Cuvelier⁴, Astrid Manders⁵, Kathleen Mar⁶, Mihaela Mircea⁷, Noelia Otero⁶, Wenche Aas⁸, Maria-Teresa Pay⁹, Valentin Raffort¹⁰, Yelva Roustan¹⁰, Mark R. Theobald¹¹, Marta G. Vivanco¹¹, Gino Brigant⁷, Andrea Cappelletti⁷, Massimo D'Isidoro⁷, Hilde Fagerli¹, Peter Wind^{1,12}

¹ Climate Modelling and Air Pollution Division, Research and Development Department, Norwegian Meteorological Institute (MET Norway), P.O. Box 43, Blindern, N-0313 Oslo, Norway

² Swedish Meteorological and Hydrological Institute, 60176 Norrköping, Sweden

³ INERIS, National Institute for Industrial Environment and Risks, Parc Technologique ALATA, 60550 Verneuil-en-Halatte, France

⁴ ex European Commission, Joint Research Centre (JRC), Ispra, Italy

⁵ TNO, Dept. Climate, Air and Sustainability, P.O. Box 80015, 3508 TA Utrecht, the Netherlands

⁶ Institute for Advanced Sustainability Studies, Potsdam, Germany

⁷ ENEA, Italian National Agency for New Technologies, Energy and Sustainable Economic Development
Via Martiri di Monte Sole 4, 40129 Bologna, Italy

⁸ Norwegian Institute for Air Research (NILU), Box 100, 2027 Kjeller, Norway

⁹ BSC, Barcelona Supercomputing Center, Centro Nacional de Supercomputación, Nexus II Building,
Jordi Girona, 29, 08034 Barcelona, Spain

¹⁰ Joint Laboratory Ecole des Ponts ParisTech - EDF R&D, CEREAs, Champs-Sur-Marne, France

¹¹ CIEMAT, Atmospheric Pollution Unit, Avda. Complutense, 22, 28040 Madrid, Spain

¹² Faculty of Science and Technology, University of Tromsø, Tromsø, Norway

Abstract: The model trend analysis for PM₁₀ and PM_{2.5}, performed within the Eurodelta-Trends experiment, covers 21 year, from 1990 through 2010, with particular focus on the period 2001-2010 for which appropriate amount of PM observations is available. Eight chemical transport models (CTM) participated in the multi-modal trend analysis: EMEP/SCS-W, CHIMERE, LOTOS-EUROS, MINNI, MATCH, WRF-Chem, CMAQ and Polyphemus (out of which six models performed trend runs for the 21-year period). The average modelled trends are somewhat smaller than the observed, though the models identify significant PM trends at more sites in the period 2001-2010. There are considerable difference in the PM trends between the regions/countries and in different seasons. Investigation of the changes in PM chemical composition during the investigated period shows that the models differ in terms of relative contribution of the individual PM components to the PM trends. For the 2001-2010 period, the effect of inter-annual meteorological variability appears more important relative to emission changes. Finally, we look at PM trends/changes during the 1990-2010 period.

Key words: PM₁₀ and PM_{2.5} trends, multi-modal assessment, observed trends.

INTRODUCTION

The Eurodelta-Trends multi-model chemistry-transport experiment was designed in order to better understand the evolution of air pollution and its drivers since the early 1990s. The main objective of the experiment is to assess the efficiency of air pollutant emissions mitigation measures in improving regional scale air quality in Europe. The presented multi-model trend analysis is a contribution to the assessment of the evolution of air pollution in the EMEP region over the 1990-2012 period coordinated by the Task Force on Monitoring and Modelling of EMEP (Cooperative Programme for Monitoring and

Evaluation of the Long-range Transmission of Air Pollutants in Europe) and presented Colette et al., 2016).

EXPERIMENT DESIGN

The set-up and input information for model runs were largely harmonized, thus allowing investigation of the uncertainties in calculation results due to the differences in model formulations. Six of the chemical transport models (CTM) participated in the multi-modal trend analysis, i.e. EMEP MSC-W, CHIMERE, LOTOS-EUROS, MINNI, MATCH, WRF-Chem, performed a 21-year run series over Europe for the 1990-2010 period. In addition, those six models and also CMAQ and Polyphemus made calculations for the years 1990, 2000 and 2010. In order to study the effect of meteorological variability, additional runs with constant 2010 emission were performed for the same years. For detailed description of the experiment see Colette et al. (2017).

The model trend analysis for PM_{10} and $PM_{2.5}$ covers 21 years, from 1990 through 2010. For the 2001-2010 period, for which enough PM observations are available from the EMEP monitoring network, the modelled trends of PM_{10} and $PM_{2.5}$ have been compared with measurement data. For both model results and observations, the Mann Kendall (MK) method has been used to identify significant trends, and Sen's slopes have been calculated to quantify PM trends.

RESULTS

Model performance for PM

Comparison with EMEP observations for the years 2001 to 2010 shows that the models underestimate PM_{10} by 14% and $PM_{2.5}$ by 11% on average. The inter-annual bias variation is mostly within 5% (up to 10%), and a certain consistency can be noticed between the models' performance (e.g. smaller underestimation of PM_{10} for 2001, 2005, 2006 and 2009 and greater underestimation for 2002, 2007, 2008, 2010). The average spatial correlation coefficients are 0.58 PM_{10} and 0.67 for $PM_{2.5}$ and in general vary only moderately between the years and the models.

Trends for 2001-2010

Figure 1 presents for the period 2001-2010 the observed and modelled changes in mean PM_{10} and $PM_{2.5}$ levels and the trend slopes at the selected set of trend-sites (Colette et al., 2016), showing the mean from 5-model ensemble and the individual models' results.




Figure 1. PM_{10} and $PM_{2.5}$ mean level change and Sen's slopes observed and according to 6-model ensemble for the period 2001-2010; bottom figures also show results from the individual models.

The PM levels from the 5-model ensemble are lower than observed, with negative biases being larger for PM₁₀ than for PM_{2.5}, but the discrepancies somewhat decrease towards the end of the period. The model calculated year-to-year PM changes follow quite well the observations, except from the end of the period. There, we see a decrease in calculated PM levels, while the measured PM increases. The resulting average modelled trends are somewhat smaller than the observed ones. For PM₁₀: calculated mean -0.31 $\mu\text{g m}^{-3} \text{ yr}^{-1}$ (or -2.0 % per yr), observed -0.42 $\mu\text{g m}^{-3} \text{ yr}^{-1}$ (or -2.3 % per yr); and for PM_{2.5}: calculated mean -0.28 $\mu\text{g m}^{-3} \text{ yr}^{-1}$ (or -2.5 % per yr), observed -0.41 $\mu\text{g m}^{-3} \text{ yr}^{-1}$ (or -3.2 % per yr). For the individual models, the correlations between annual mean modelled and measured PM₁₀ are: 0.85 for CHIM, 0.88 for EMEP, 0.62 for LOTO, 0.80 for MINNI and 0.84 for MATCH. For PM_{2.5}, the correlations between the model results and observations are even higher, with the corresponding values being 0.95, 0.94, 0.83, 0.90 and 0.95.

The calculations and observations show considerable difference in regional PM trends. Figure 2 shows the mean values of Sen's slope for the period of 2001-2010 calculated by the ensemble of six models and measured at EMEP sites. The model mean and observed annual levels of PM₁₀ and PM_{2.5} were in general decreasing overall in Europe during the 2001-2010. Compared to observations, the models identified significant trends in PM₁₀ and PM_{2.5} for larger geographical regions and more locations (though underestimating the trend slopes).




Figure 2. Mean Sen's slopes for PM₁₀ and PM_{2.5} trends in 2001-2010 ($\mu\text{g m}^{-3} \text{ year}^{-1}$) calculated by 6-model ensemble (described in Colette et al. (2016), Appendix A3.) Shown are modelled trends – coloured contours (grey means insignificant trends) and observed trends - coloured triangles (significant) and circles (insignificant).

The study reveals quite a large variability of observed mean trends at the individual sites, ranging from -0.93 to 0.003 $\mu\text{g m}^{-3} \text{ yr}^{-1}$ (from -0.83 to -0.308 $\mu\text{g m}^{-3} \text{ yr}^{-1}$ for significant trends). Due to large meteorological variability and the relatively short studied period, the observations do not indicate significant PM₁₀ trends at 14 out of 29 sites; and only at 8 out of 18 PM_{2.5} trend-sites, significant trends were observed. The model calculated mean trends at different sites vary less compared to the observations between the sites. In general, the models tend to under-predict significant observed trends, whereas the modelled trend are closer to, or overestimate observations at the sites with insignificant measured trends.




Figure 3. Seasonal mean concentrations of PM₁₀ averaged over 29 (blue) and PM_{2.5} averaged over 18 sites (red). Solid lines - observations, dotted lines – 5-model ensemble; grey lines – Sen's slopes.

Analysis of seasonal profiles shows considerable variability, with the largest trends observed and calculated for summer and the smallest for winter (Figure 3).

The models agree that the chemical composition of European PM₁₀ was changing in the course of 2001-2010 decade, with NO₃ becoming increasingly more important constituent compared to SO₄, NH₄ and carbonaceous particles from primary PM. However, the models differ in terms of relative contribution of the individual PM components to the PM trends.

Trends for 1990-2010

PM trends for the period of 1990-2010 could only be studied based on model calculations, which show downward trends over the whole of Europe. The mean estimated trends are $-0.37 \mu\text{g m}^{-3} \text{yr}^{-1}$ for PM₁₀ and $-0.32 \mu\text{g m}^{-3} \text{yr}^{-1}$ for PM_{2.5}, though the slope values vary considerably between the countries. All models calculate a larger decrease of PM levels from 1990 to 2000 compared to the 2000-2010 period, except from Northern Europe (as example from the EMEP model in Figure 4).




Figure 4. Mean PM₁₀ trends ($\mu\text{g m}^{-3} \text{yr}^{-1}$) for the individual countries for 1990-2000 and 2000-2010 periods (calculated with the EMEP model).

Finally, our study shows considerable effect of the inter-annual changes in meteorological conditions on the modelled 1990-2010 trends, and for the shorter period of 2001-2010 with smaller emission changes the meteorology appears more important than the emission changes.

REFERENCES

Colette, A., Andersson, C., Manders, A., Mar, K., Mircea, M., Pay, M.-T., Raffort, V., Tsyro, S., Cuvelier, C., Adani, M., Bessagnet, B., Bergström, R., Briganti, G., Butler, T., Cappelletti, A., Couvidat, F., D'Isidoro, M., Doumbia, T., Fagerli, H., Granier, C., Heyes, C., Klimont, Z., Ojha, N., Otero, N., Schaap, M., Sindelarova, K., Stegehuis, A. I., Roustan, Y., Vautard, R., van Meijgaard, E., Vivanco, M. G., and Wind, P., 2017: EURODELTA-Trends, a multi-model experiment of air quality hindcast in Europe over 1990–2010, *Geosci. Model Dev.*, **10**, 3255–3276, <https://doi.org/10.5194/gmd-10-3255-2017>.

Colette, A., Aas, W., et al.: 2016: Air pollution trends in the EMEP region between 1990 and 2012. Joint Report of: EMEP Task Force on Measurements and Modelling (TFMM), Chemical Coordinating Centre (CCC), Meteorological Synthesizing Centre-East (MSC-E), Meteorological Synthesizing Centre-West (MSC-W). EMEP/CCC-Report 1/2016. Norwegian Institute for Air Research P.O. Box 100, NO-2027 Kjeller, Norway.