detritus | Multidisciplinary Journal for Waste Resources & Residues

CISA

# PROPOSITION OF THRESHOLD FOR WASTE CONTAMINATED WITH MERCURY (COMPOUNDS) IN APPLICATION OF THE MINAMATA CONVENTION ON MERCURY AND IMPACT ASSESSMENT

Pierre Hennebert *

*INERIS (French National Institute for Industrial Environment and Risks), BP 2, F-60550 Verneuil-en-Halatte, France*

**Article Info:**
Received:
12 March 2019
Revised:
18 June 2019
Accepted:
20 June 2019
Available online:
28 June 2019

**Keywords:**
Concentration limit
GHS
Ecotoxicity
Environmentally sound management

**ABSTRACT**

The Minamata Convention on Mercury is a global treaty to protect human health and the environment from the adverse effects of mercury. Environmentally sound management of waste is under discussion. This note proposes a threshold for waste of category c) *Contaminated with mercury or mercury compounds* to be disposed of (Article 11 of the Convention), using the Globally Harmonized System of classification and labelling of chemicals of the United Nations (GHS - UNEP, 2017). Mercury and mercury compounds are classified as substances for the physical, health and environmental hazards categories. The thresholds of mercury and mercury compounds classifying a mixture as hazardous for the different hazard categories (physical, health, environmental) are "Presence", >0.1% and >0.0025% (25 mg mercury/kg of waste) respectively. For impact assessment, this threshold is then compared with large data set of hazardous (793 data), potentially hazardous (depending on the concentration of hazardous substances) (55 data), as well as natural or non-polluted anthropized media (composts, sediments, agricultural soils) (21 784 data) from France. This demonstrates that 75% of the hazardous waste have higher total mercury concentration, that potentially hazardous waste samples have lower concentrations, and that all composts, agricultural soils and marine sediments and 99% of the fluvial sediments have lower concentrations. So, this threshold will not classify common industrial waste or natural media as requiring special treatment for safe disposal, but well a large part of industrial hazardous waste.

## 1. INTRODUCTION

The Minamata Convention on Mercury (UNEP 2017) is a global treaty to protect human health and the environment from the adverse effects of mercury. Regular conference of the parties progress in technical recommendations to "make mercury history". Environmentally sound management of waste is one point under discussion.

The Convention defines in Article 11 "Mercury wastes":

*"…2. For the purposes of this Convention, mercury wastes means substances or objects:*
*(a) Consisting of mercury or mercury compounds;*
*(b) Containing mercury or mercury compounds; or*
*(c) Contaminated with mercury or mercury compounds,*
*in a quantity above the relevant thresholds defined by the Conference of the Parties, in collaboration with the relevant bodies of the Basel Convention in a harmonized manner, that are disposed of or are intended to be disposed of or are required to be disposed of by the provisions of national law or this Convention. This definition excludes overburden, waste rock and tailings from mining, except from primary mercury mining, unless they contain mercury or mercury compounds above thresholds defined by the Conference of the Parties…"*

The scope of the paper is to provide a reliable concentration limit to the Minamata Convention. This paper uses the Globally Harmonized System (GHS) of classification and labelling of chemicals of the United Nations to propose a threshold and compare it with concentrations observed in waste and natural media. The method is explained in detail in the paper, with a focus on ecotoxicity, which appears to be the property with the lowest dangerous ranking.

In this paper, the "concentration limit" used in the GHS is the equivalent of "threshold" of the Minamata Convention on Mercury, and "substance" used in the GHS is the equivalent of "compound" of the Minamata Convention on Mercury.

* Corresponding author:
Pierre Hennebert
email: pierre.hennebert@ineris.fr

Detritus / Volume 06 - 2019 / pages 25-31
https://doi.org/10.31025/2611-4135/2019.13822
© 2019 Cisa Publisher. Open access article under CC BY-NC-ND license

## 2. MATERIAL AND METHODS

### 2.1 Properties of mercury and mercury substances in the GHS

It is proposed to use the Globally Harmonized System (GHS) of classification and labelling of chemicals of the United Nations (last version: UNEP, 2017). The European Union has adopted the GHS system in 2008 (CLP, 2008) and has developed an official list of the hazard properties of 4 249 substances, as well as a registration and self-classification system for producers and importers of chemical products (REACh system). This list is an Annex of the CLP regulation and can be downloaded in a spreadsheet template (CLP, 2018) and is used here. These "harmonized" (at EU level) data were built by working group of experts and have been used to collect consistent data on mercury and mercury compounds and to propose threshold consistent with the regulation.

### 2.2 Particular case: ecotoxicity in the GHS

The different categories of ecotoxicity are attributed to substances from experimental laboratory ecotoxicological standardized tests. Organisms are submitted to different concentration of the substance in their living medium, and the concentration producing 50% (or x %) of mortality or effect after a given time (called $LC_{50}$ or $EC_{50}$ or $EC_x$ depending on test), or the highest concentration producing no observed effect (called NOEC) is measured. These concentrations expressed in mg of substance per liter of living medium are compared to concentrations of Table 1, and a category is attributed to the substance.

For substances with experimental results < 1 mg/l, the GHS uses multiplying factors called "M factors" to fine tune the classification of substances and mixtures with these substances. M factors are determined by the lowest $L(E)C_{50}$ and NOEC experimental values (Table 2). Mercury and mercury compounds are non-rapidly degradable compounds.

Experimental ecotoxicity data of mercury and mercuric substances were taken from a reference UE publication (UE 2005), to determine M-factors for mercury and mercury compounds.

The ecotoxicity of mixtures with ecotoxic substances at a given concentration can be assessed by the calculation rules presented at Table 3 (copy of Table 4.1.3 of the GHS).

The European Union updated criteria on Ecotoxicity classification of waste (Regulation 997/2017, EU 2017). For simplicity of classification of waste with most of the time unknown mineral substances composition, this regulation does not use the multiplying factors M-factors (that are attributes of substances). M-factors were developed in the GHS to fine tune the ecotoxicity of ecotoxic substances ($EC_{50}$ or NOEC < 1 mg/L), and to avoid the creation of multiple hazard statement codes to reflect the different grades of ecotoxicity. The resulting concentration limit for waste by calculation method as proposed by the 997/2017 Regulation for chronic ecotoxicity for mercury and mercury compounds is 0.25% or 2500 mg/kg, hundred times higher than the concentration calculated here (see Results). Many experts believe that the abandonment of M factors for waste is irrelevant. This method has not been used here.

### 2.3 Data of concentration of mercury in waste and natural media

Data of hazardous waste, potentially hazardous waste

**TABLE 1:** Categories for substances hazardous to the aquatic environment for short-term (acute) and long-term (chronic) aquatic hazard (extract of Table 4.1.1 of GHS).

(a) Short-term (acute) aquatic hazard
- Category Acute 1: $L(E)C_{50}$ fish, crustacea, algae ≤ 1 mg/l
- Category Acute 2: $L(E)C_{50}$ fish, crustacea, algae > 1 but ≤ 10 mg/l
- Category Acute 3: $L(E)C_{50}$ fish, crustacea, algae > 10 but ≤ 100 mg/l

(b) Long-term (chronic) aquatic hazard

(i) Non-rapidly degradable substances for which there are adequate chronic toxicity data available
- Category Chronic 1: Chronic NOEC or $EC_x$ fish, crustacea, algae ≤ 0.1 mg/l
- Category Chronic 2: Chronic NOEC or $EC_x$ fish, crustacea, algae ≤ 1 mg/l

(ii) Rapidly degradable substances for which there are adequate chronic toxicity data available
- Category Chronic 1: Chronic NOEC or $EC_x$ fish, crustacea, algae ≤ 0.01 mg/l
- Category Chronic 2: Chronic NOEC or $EC_x$ fish, crustacea, algae ≤ 0.1 mg/l
- Category Chronic 3: Chronic NOEC or $EC_x$ fish, crustacea, algae ≤ 1 mg/l

**TABLE 2:** Multiplying factors (M factors) for substances for highly toxic ingredients or mixture (categories Acute 1 and Chronic 1) (Table 4.1.5 of GHS).

| Acute toxicity | M factor | Chronic toxicity | M factor | |
|---|---|---|---|---|
| $L(E)C_{50}$ value (mg/l) | | NOEC value (mg/l) | Non-rapidly degradable ingredients | Rapidly degradable ingredients |
| 0.1 < $CL(E)_{50}$ ≤ 1 | 1 | 0.01 < NOEC ≤ 0.1 | 1 | - |
| 0.01 < $CL(E)_{50}$ ≤ 0.1 | 10 | 0.001 < NOEC ≤ 0.01 | 10 | 1 |
| 0.001 < $CL(E)_{50}$ ≤ 0.01 | 100 | 0.0001 < NOEC ≤ 0.001 | 100 | 10 |
| 0.0001 < $CL(E)_{50}$ ≤ 0.001 | 1000 | 0.00001 < NOEC ≤ 0.0001 | 1000 | 100 |
| 0.00001 < $CL(E)_{50}$ ≤ 0.0001 | 10000 | 0.000001 < NOEC ≤ 0.00001 | 10000 | 1000 |
| (continue in factor 10 intervals) | | (continue in factor 10 intervals) | | |

and non-hazardous waste and natural media have been gathered:

- The concentrations of mercury, cadmium and lead in 793 hazardous waste (data from a hazardous waste management company) (Hennebert, 2012);
- The 55 available data concentrations from an INERIS database for the following waste: car shredding residue (11), epoxy powder (1), excavated soil underlying a road (3), excavated soil (11), foundry sand (1), municipal solid waste incinerator bottom ash - MSWI BA (20) , paint residue (1), phosphogypsum (2), sand blasting residue (1) and sand from incineration fluidized bed (4);
- The heavy metals in different composts (379 samples) from organic fraction of municipal wastes (separately collected or mechanically sorted) of 30 sites in France (Zdanevitch 2012);
- The results of routine quality monitoring of sediments (11 791 samples of fluvial sediments and 816 samples of marine sediments) of France by the Water Agencies (Padox and Hennebert 2010a, b);
- The data from routine analysis of agricultural soils (8 798 samples) are gathered at the French level by soil scientists (Gissol, 2018).

## 3. RESULTS AND DISCUSSION

### 3.1 Properties of mercury and mercury substances in the GHS

The mercury and nine mercury substances from the list are presented at Table 6. Two "generic entries" are also listed. Most frequently in waste, field measurement with fluorimeter and routine laboratory analysis will deliver total mercury concentrations (metallic or not) rather than mercury substances concentrations. If the exact mercury substances present in the waste is not known, which is frequently the case in waste, these "generic entries" are used in the EU (line 2 and 3 of the substances list in Table 6: "inorganic compounds of mercury with the exception of mercuric sulphide and those specified elsewhere in this Annex" and "organic compounds of mercury with the exception of those specified elsewhere in this Annex"). In total twelve entries are used for mercury and mercury substances, with their hazard statement codes (Hxxx).

Mercury and mercury substances are all classified Acute Toxic when the three routes (oral, dermal and inhalation) are considered, classified Specific Target Organ Toxic Single Exposure (1 substance: calomel) or Repeated Exposure (11 other substances), Ecotoxic Aquatic Acute Level 1 and Ecotoxic Aquatic Chronic Level 1. Additionally, mercury dichloride is classified mutagenic and reprotoxic, and elemental mercury is reprotoxic. Mercury difulminates and dimercury dicyanide dioxide are explosive (Table 6).

#### *3.1.1 Classification of mercury and mercury substances (GHS, EU data) for ecotoxicity*

Experimental ecotoxicity data of mercury and mercuric substances from a reference UE publication (UE 2005) are presented at Table 4. A summary can be found on the INERIS portal (INERIS 2018). The resulting M factors are

**TABLE 3:** Classification of a mixture for short-term (acute) and long-term (chronic) hazard based on summation of the concentrations of classified ingredients (Tables 4.1.3 and 4.1.4 of GHS).

| Category | Sum of the concentrations (in %) of ingredients classified as: | Mixture is classified as: |
|---|---|---|
| Short-term (acute) hazard | Acute 1 * M ≥ 25% | Acute 1 |
| | (M*10*Acute 1) + Acute 2 ≥ 25% | Acute 2 |
| | (M*100*Acute 1) + (10*Acute 2) + Acute 3 ≥ 25% | Acute 3 |
| Long-term (acute) hazard | Chronic 1 * M ≥ 25% | Chronic 1 |
| | (M*10*Chronic 1) + Chronic 2 ≥ 25% | Chronic 2 |
| | (M*100*Chronic 1) + (10*Chronic 2) + Chronic 3 ≥ 25% | Chronic 3 |
| | Chronic 1 + Chronic 2 + Chronic 3 + Chronic 4 ≥ 25% | Chronic 4 |

**TABLE 4:** Experimental $L(E)C_{50}$ and NOEC of mercury and mercury substances (UE 2005) and corresponding ecotoxicity level and multiplying factors (M factors).

| Water | | Fresh | | | Marine | | Classification | |
|---|---|---|---|---|---|---|---|---|
| Ecotoxicity Organisms | Tests results (mg/L) | Ecotoxicity category (Table 2) | M-factor (Table 3) | Tests results (mg/L) | Ecotoxicity category (Table 2) | M-factor (Table 3) | Category | Proposed M factor |
| Acute | $L(E)C_{50}$ | | | $L(E)C_{50}$ | | | Acute 1 | 100 |
| Algae | 0.010000 | Acute 1 | 100 | 0.010000 | Acute 1 | 100 | | |
| Invertebrate | 0.010000 | Acute 1 | 100 | 0.003500 | Acute 1 | 100 | | |
| Fish | 0.000700 | Acute 1 | 1000 | 0.070000 | Acute 1 | 10 | | |
| Chronic | NOEC | | | NOEC | | | Chronic 1 | 100 |
| Algae | 0.000200 | Chronic 1 | 100 | 0.000900 | Chronic 1 | 100 | | |
| Invertebrate | 0.000290 | Chronic 1 | 100 | 0.000100 | Chronic 1 | 1000 | | |
| Fish | 0.000620 | Chronic 1 | 100 | 0.005000 | Chronic 1 | 10 | | |

most of the time 100 (with two values of 10 and two values of 1000 in a set of 12 values). It is proposed here to use M = 100 for acute and chronic ecotoxicity.

Using the calculation rules of Table 3 and the M factors obtained in Table 4, the concentration limits classifying a mixture containing mercury or mercury substances as ecotoxic acute and chronic are presented at Table 5. For full classification of mixtures, all the other ecotoxic substances must also be used, but this is not the question here. The level 3 has the lowest concentration: a mixture is hazardous if the concentration of mercury and/or mercury substances is greater than 0.0025% (mass/mass), or 25 mg/kg (sometimes expressed as 25 parts per million - ppm). These concentrations are reported as concentration limit for Environmental hazard in Table 6.

*3.1.2 Proposed concentration limits for waste containing mercury or mercury compounds for physical, health and environmental hazard classification*

The concentration limits are presented in the last row of Table 6, with the lowest by category of hazard in color. For human acute toxicity, concentration limits are derived from acute toxicity estimates for mixtures of the GHS. Some substances have specific concentration limits for some hazard class category in the EU list of substances, but they were not used here. For physical hazard, the requirement is "presence". For health hazards, the lowest concentration is 0.1% of mercury or mercury substances in the mixture (Specific target organ toxicity, repeated exposure). For environmental hazard, the concentration limit is 0.0025% or 25 mg/kg.

### 3.2 Impact assessment of proposed threshold

Numerous data (> 22 000) from France, for different waste streams were collected to perform impact assessment of the proposed threshold:

- Hazardous waste (according to the EU List of Waste);
- Potentially hazardous waste (depending on the concentration of hazardous substances in it) (so-called "mirror entries" in the EU List of Waste), including municipal solid waste incinerator (MSWI) bottom ash;
- Composts, sediments and agricultural soils.

Regulatory considerations on municipal sewage sludge (biosolids) and waste intended to be used as fertilizers are also presented.

More than 75% of the hazardous wastes have more than 25 mg Hg/kg. Data of Cd and Pb are presented, to illustrate that other hazardous elements are most of the time also present in these wastes (Table 7).

The Hg concentration for 55 potentially hazardous waste are presented at Figure 1 (left) and for MSWI BA at Figure 1 (right).

All the reported concentrations are lower than 25 mg/kg. For the MSWI BA, 8 samples have concentrations lower than the quantification limits of the laboratory (between 0.175 mg/kg and 5 mg/kg).

The distributions of Hg in different composts from organic fraction of municipal wastes (separately collected or mechanically sorted) of 30 sites in France (Zdanevitch, 2012) are presented in Table 8.

The results of routine quality monitoring of sediments are joined to the table. The network aims to monitor pollution and hot spots are more intensively sampled. The regulatory concentration limits of Hg for these sediments for reuse in natural environment are 1 mg/kg and 0.8 mg/kg respectively. One percent of the fluvial sediments should be specifically managed for mercury, with the threshold of 25 mg/kg, and no marine sediment. These samples are contaminated by other heavy-metals (As, Cu, Cd, Pb, Zn - result not shown) and are in all the case identified in survey and separated during dredging.

Data from routine analysis of agricultural soils are presented (Gissol 2018). A detailed analysis is available (Baize et al 2007). Another publication deals among others with local geological "anomaly" of heavy metals (Baize 2000). According to the authors, the higher concentrations originates from the parent material of the soil (most of the case) or from industrial inputs. All the concentrations are lower than 25 mg/kg (Table 8).

For municipal sewage sludge, the EEC Council Directive on the protection of the environment, and in particular of the soil, when sewage sludge is used in agriculture (EEC 1986) (updated) states in its Annex I B the limit values for heavy-metal concentrations in sludge for use in agriculture for mercury: 16 to 25 mg/kg of dry matter. In France, the corresponding decree sets a concentration limit of 10 mg/kg of dry matter (RF 1998).

For reuse of waste as fertilizing products in the Circular Economy package (EC 2016), the EC has confirmed the concentrations of Hg in fertilizers, culture medium and

**TABLE 5:** Classification of a mixture for acute and chronic aquatic ecotoxicity based on summation of the concentrations of classified ingredients (source: UNEP 2017), containing mercury or mercury substances (M factor Acute and M factor Chronic = 100). In green: the lowest concentration limit.

| Level of ecotoxicity | Ecotoxic Acute if sum of | Ecotoxic Acute (M=100) if sum of mercury and mercury substances | Ecotoxic Chronic if sum of | Ecotoxic Chronic (M=100) if sum of mercury and mercury substances |
|---|---|---|---|---|
| 1 | Acute 1 * M ≥25% | ≥ 0.25% | Chronic 1 * M ≥25% | ≥ 0.25% |
| 2 | (M*10*Acute 1) + Acute 2 ≥ 25% | ≥ 0.025% | (M*10*Chronic 1) + Chronic 2 ≥ 25% | ≥ 0.025% |
| 3 | (M*100*Acute 1) + (10*Acute 2) + Acute 3 ≥ 25% | ≥ 0.0025% = 25 mg/kg | (M*100*Chronic 1) + (10*Chronic 2) + Chronic 3 ≥ 25% | ≥ 0.0025% = 25 mg/kg |
| 4 | - | - | Chronic 1 + Chronic 2 + Chronic 3 + Chronic 4 ≥ 25% | ≥ 25% |

**TABLE 6:** Classification of mercury and mercury substances with the GHS in the EU, and corresponding concentration limits classifying a mixture containing mercury or mercury substances as hazardous. In color: the lowest concentration limits for physical (brown), health (blue) and environmental (green) hazards.

| Hazard group | | Physical | | Health | | | | | | | | | | | | | | | Environmental | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Hazard Property | | Explosive | | Acute toxicity | | | | | | | Irritant | | | STOT* | | | Muta-genicity | Repro-ductive toxicity | | Acute M=100 | Chronic M=100 |
| | | | | Oral | | | Dermal | | Inhal. | | | | | SE* | RE* | | | | | | |
| Substances and hazard statement codes | CAS | H200 | H201 | H300 Cat 2 | H301 | H302 | H310 Cat 1 | H311 | H330 Cat 2 | H331 | H314 | H315 | H319 | H335 | H372 | H373 | H341 | H360 | H361 | Cat 1 H400 | Cat 1 H410 |
| Mercury | 7439-97-6 | | | | | | | | X | | | | | | X | | | X | | X | X |
| Inorganic compounds of mercury with the exception of mercuric sulphide and those specified elsewhere in this Annex | - | | | X | | | X | | X | | | | | | | X | | | | X | X |
| Organic compounds of mercury with the exception of those specified elsewhere in this Annex | - | | | X | | | X | | X | | | | | | | X | | | | X | X |
| Dimethylmercury [1]; diethylmercury [2] | 593-74-8 [1]; 627-44-1 [2] | | | X | | | X | | X | | | | | | | X | | | | X | X |
| Dimercury dicyanide oxide; mercuric oxycyanide | 1335-31-5 | | X | | X | | | X | | X | | | | | | X | | | | X | X |
| Dimercury dichloride; mercurous chloride; calomel | 10112-91-1 | | | | | X | | | | | | X | X | X | | | | | | X | X |
| Mercury dichloride; mercuric chloride | 7487-94-7 | | | X | | | | | | | X | | | | X | | X | | X | X | X |
| 2-methoxyethylmercury chloride | 123-88-6 | | | | X | | | | | | X | | | | X | | | | | X | X |
| Phenylmercury acetate | 62-38-4 | | | | X | | | | | | X | | | | X | | | | | X | X |
| Phenylmercury nitrate [1]; phenylmercury hydroxide [2]; basic phenylmercury nitrate [3] | 55-68-5 [1]; 100-57-2 [2]; 8003-05-2 [3] | | | | X | | | | | | X | | | | X | | | | | X | X |
| Mercury difulminate; mercuric fulminate; fulminate of mercury | 628-86-4 | X | | | X | | | X | | X | | | | | | X | | | | X | X |
| Mercury difulminate; mercuric fulminate; fulminate of mercury [≥ 20 % phlegmatiser] | 628-86-4 | | X | | X | | | X | | X | | | | | | X | | | | X | X |
| Concentration limits = tresholds | Hazardous if | Presence | Presence | ≥ 0.25% | ≥ 5% | ≥ 25% | ≥ 0.25% | ≥ 15% | ≥ 0.5% | ≥ 10% | ≥ 1% | ≥ 10% | ≥ 10% | ≥ 20% | ≥ 1% | ≥ 0.1% lowest | ≥ 1% | ≥ 0.3% | ≥ 3% | ≥ 0.0025% | ≥ 0.0025% |

** STOT SE = specific target organ toxicity – single exposure, STOT RE = specific target organ toxicity – repeated exposure*

**TABLE 7:** Distribution of Hg (in red: > proposed threshold of 25 mg Hg/kg), Cd and Pb in hazardous waste in France.

| Element | Hg | Cd | Pb |
|---|---|---|---|
| Number of data | 793 | 2266 | 2856 |
| Mean | 7121 | 1272 | 6961 |
| Min | 0.1 | 0.0 | 1.1 |
| 1% | 0.3 | 0.5 | 6.3 |
| 5% | 10 | 2 | 40 |
| 10% | 10 | 4 | 69 |
| 25% | 190 | 14 | 162 |
| 50% = median | 836 | 55 | 1758 |
| 75% | 5336 | 129 | 3464 |
| 90% | 12228 | 394 | 8907 |
| 95% | 16835 | 2481 | 29192 |
| 99% | 121368* | 9725 | 101794* |
| Max | 544018* | 728708* | 550423* |

** Probable analytical bias (X ray fluorescence analysis)*

**TABLE 8:** Concentration of Hg (mg/kg) in compost, fluvial and marine sediments, and agricultural soils (21 784 data) (in red: > proposed threshold of 25 mg Hg/kg).

| Element | Compost | Fluvial sediments | Marine sediments | Agricultural soils |
|---|---|---|---|---|
| N samples | 379 | 11 791 | 816 | 8798 |
| Mean | 0.39 | 1.22 | 0.50 | 0.08 |
| Min | 0.00 | 0.002 | 0.01 | 0.01 |
| 1% | 0.02 | 0.005 | 0.02 | |
| 10% | 0.09 | 0.03 | 0.05 | 0.02 |
| 25% | 0.20 | 0.05 | 0.08 | 0.03 |
| 50% = median | 0.30 | 0.13 | 0.13 | 0.05 |
| 75% | 0.50 | 0.40 | 0.26 | 0.07 |
| 90% | 0.80 | 1.30 | 0.53 | 0.11 |
| 99% | 1.68 | 31.0 | 2.47 | |
| Max | 2.40 | 200 | 112 | 11.6 |

soils improvers of 1 mg/kg and in one case 2 mg/kg, that are in the Regulation relating to fertilizers (EC 2003).

## 4. CONCLUSION

The Globally Harmonized System of the UN classifies mixtures with concentration of mercury and mercury substances as hazardous (aquatic acute and chronic ecotoxicity of level 3) if their concentration is greater than or equal to 0.0025%, or 25 mg/kg. This concentration limit could be used as threshold for disposal of waste of category “c” according to the Minamata Convention.

The impact assessment shows that this concentration will not classify any of the common industrial waste or composts, sediments of soils to be managed specifically for mercury, but well the hazardous waste, that are already stabilized or solidified before landfilling in special landfills for hazardous waste, and very few (1%) contaminated fluvial sediments (that in all the case are also contaminated with other heavy metals), according to French data. It is also consistent with the present concentration limits set in the EU for reuse of municipal sewage sludge and other waste as fertilizing products.

## REFERENCES


Baize D, Saby N, Deslais W. 2007. Teneurs en huit éléments en traces (Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn) dans les sols agricoles en France. Résultats d'une collecte de données à l'échelon national. In French. INRA - Orléans Unités Science du sol et InfoSol. 26 janvier 2007. http://www.gissol.fr/programme/bdetm/_rapport_anademe/rapport/03.php#3-2-2

Baize D. 2000. Teneurs totales en "métaux lourds" dans les sols français - résultats généraux du programme ASPITET. In French. Le Courrier de l'environnement n°39, février 2000. http://www.inra.fr/dpenv/baizec39.htm

CLP 2018. Annex A spreadsheet file of EU classified substances https://echa.europa.eu/information-on-chemicals/annex-vi-to-clp

EC 1986. Council Directive of 12 June 1986 on the protection of the environment, and in particular of the soil, when sewage sludge is used in agriculture (86/278/EEC) (OJ L 181, 4.7.1986, p. 6) https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:01986L0278-20090420&qid=1527858946053&from=EN

EC 2003. REGULATION (EC) No 2003/2003 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 13 October 2003 relating to fertilisers. https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32003R2003&rid=1

EC 2008 (CLP). Regulation EC No 1272/2008 of the European Parliament and of the Council of 16 December 2008 on classification, labelling and packaging of substances and mixtures, amending and repealing Directives 67/548/EEC and 1999/45/EC, and amending Regulation (EC) No 1907/2006. 31.12.2008 Official Journal of the European Union L 353/1. http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008R1272&rid=1






**FIGURE 1:** Distribution of total Hg concentration in 55 potentially hazardous waste (left), among which 20 samples of Municipal Solid Waste Incinerator Bottom Ash (right).

EC 2016. Circular Economy Package - Proposal for a Regulation of the European Parliament and of the COuncil laying down rules on the making available on the market of CE marked fertilising products and amending Regulations (EC) No 1069/2009 and (EC) No 1107/2009. https://eur-lex.europa.eu/resource.html?uri=cellar:d117e80d-ec28-11e5-8a81-01aa75ed71a1.0018.02/DOC_1&format=PDF

EEC 1986. Directive of 12 June 1986 on the protection of the environment, and in particular of the soil, when sewage sludge is used in agriculture (86/278/EEC) (OJ L 181, 4.7.1986, p. 6) https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:01986L0278-20090420&qid=1527858946053&from=EN

EU 2005 Environmental Quality Standards (EQS), Substance data sheet for priority substance N°21: Mercury and its compounds, CAS-N°7439-97-6, 21 p. https://circabc.europa.eu/sd/a/ff8e163c-71f6-4fc0-98ef-875a20add4c8/21_Mercury_EQSdatasheet_150105.pdf

EU 2017. Council Regulation (EU) 2017/997 of 8 June 2017 amending Annex III to Directive 2008/98/EC of the European Parliament and of the Council as regards the hazardous property HP 14 'Ecotoxic'. Official Journal of the European Union. 14.6.2017. L 150/1. https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32017R0997&rid=1

Gissol 2018. http://www.gissol.fr/programme/bdetm/_rapport_anademe/rapport/03.php#3-2-2 (consulted Dec. 2018).

Hennebert P. 2012. Classification of industrial waste for hazard properties HP4, HP6, HP8, HP13 and HP14 criteria based on substance concentrations, and impact assessment of options for HP14 on classification of various wastes, composts, sediments and soils. Technical Report INERIS DRC-12-125740-03014A. 10/03/2012. 104 p. Available upon request (pierre.hennebert@ineris.fr)

INERIS 2018. https://substances.ineris.fr/fr/substance/1183 (consulted Dec. 2018).

Padox J-M, Hennebert P. 2010a. Qualité chimique des sédiments fluviaux en France : Synthèse des bases de données disponibles. In French. Rapport d'étude INERIS-DRC-10-105335-04971A. 03/06/2010. 101 p. https://www.ineris.fr/sites/ineris.fr/files/contribution/Documents/r-10-04971a-action22.pdf

Padox J-M, Hennebert P. 2010b. Qualité chimique des sédiments marins en France : Synthèse des bases de données disponibles. In French. Rapport d'étude INERIS-DRC-10-105335-11618A. 12/08/2010. 103 p. https://www.ineris.fr/fr/qualite-chimique-des-sediments-marins-en-france-synthese-des-bases-de-donnees-disponibles

RF 1998. Arrêté du 8 janvier 1998 fixant les prescriptions techniques applicables aux épandages de bous sur les sols agricoles pris en application du décret n° 97-1133 du 8 décembre 1997 relatif à l'épandage des boues issues du traitement des eaux usées. In French. https://www.legifrance.gouv.fr/jo_pdf.do?id=JORFTEXT000000570287&pageCourante=01563

UN 2017 (GHS). Globally Harmonized System of classification and labelling of chemicals (GHS). Seventh revised edition. 534 p. http://www.unece.org/fileadmin/DAM/trans/danger/publi/ghs/ghs_rev07/English/ST_SG_AC10_30_Rev7e.pdf

UNEP 2017. Minamata Convention on Mercury – Text and Annexes. 71 p. http://mercuryconvention.org/Portals/11/documents/Booklets/COP1%20version/Minamata-Convention-booklet-eng-full.pdf

Zdanevitch I. 2012. Etude comparative de la qualité de composts et de digestats issus de la fraction fermentescible d'ordures ménagères collectée séparément ou en mélange. In French. Report 24/02/2012 N° INERIS DRC-12-125976-02525A. 92 pp. https://www.ineris.fr/sites/ineris.fr/files/contribution/Documents/ineris-drc-12-125976-02525-b-red-1337010661.pdf