



Comparing Sentinel-5P TROPOMI NO₂ column observations with the CAMS-regional air quality ensemble

Henk Eskes¹, John Douros¹, Jos van Geffen¹, Folkert Boersma¹, Steven Compernelle², Gaia Pinardi², Anne Blechschmidt³, Vincent-Henri Peuch⁴, Augustin Colette⁵, and Pepijn Veefkind¹

¹KNMI, R&D Satellite Observations, De Bilt, Netherlands (eskes@knmi.nl)

²Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Ringlaan 3, 1180 Uccle, Belgium

³Institute of Environmental Physics, University of Bremen, IUP-UB, Bremen, Germany

⁴European Centre for Medium-range Weather Forecast (ECMWF), Shinfield Park, Reading, UK

⁵National Institute for Industrial Environment and Risks (INERIS), Verneuil-en-Halatte, France

The Sentinel-5P TROPOMI instrument provides unique observations of atmospheric trace gases at a high resolution of about 5 km with near-daily global coverage, resolving individual sources like thermal power plants, industrial complexes, fires, medium-scale towns, roads and shipping routes. These datasets are especially well suited to test high-resolution regional-scale air quality (AQ) models and provide valuable input for regional emission inversion systems. In Europe, the Copernicus Atmosphere Monitoring Service (CAMS) has implemented an operational regional AQ forecasting capability for Europe based on an ensemble of 7 up to 11 European models. In the presentation we show comparisons between TROPOMI observations of nitrogen dioxide (NO₂) and the CAMS AQ forecasts and analyses of NO₂. We discuss the different ways of making these comparisons, and present the quantitative results for time series for regions and cities between May 2018 to March 2021, for summer and winter months and individual days. We demonstrate the importance of the free tropospheric contribution to the tropospheric column, and include profiles from the CAMS configuration of the ECMWF's global integrated model above 3 km altitude in the comparison. The models generally capture the fine-scale daily and averaged features observed by TROPOMI in much detail. In summer, the quantitative comparison of the NO₂ tropospheric column shows a close agreement, but in winter we find a significant discrepancy in the average column amount over Europe. Recently a new TROPOMI NO₂ reprocessing with processor version 2.3.1 has become available, and impact of this new version on the comparisons is discussed.

As spin-off, we present a new TROPOMI NO₂ level-2 data product for Europe, based on the replacement of the original TM5-MP generated global a priori profile (1x1 degree resolution) by the regional CAMS ensemble profile at 0.1x0.1 degree resolution. This a-priori replacement leads to significant changes in the TROPOMI retrieved tropospheric column, with typical increases at the emission hotspots in the order of 20%.

The European NO₂ product is compared with ground-based remote sensing measurements of 6

PANDORA instruments of the Pandora global network and 8 MAX-DOAS instruments. As compared to the standard S5P tropospheric NO₂ column data, the overall bias of the new product is smaller owing to a reduction of the multiplicative bias linked to the profile shape.